

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
31 December 2003 (31.12.2003)

PCT

(10) International Publication Number  
**WO 2004/000871 A2**

(51) International Patent Classification<sup>7</sup>: C07K

(21) International Application Number:  
PCT/US2003/019441

(22) International Filing Date: 20 June 2003 (20.06.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/390,185 21 June 2002 (21.06.2002) US

(71) Applicant (for all designated States except US): MON-  
SANTO TECHNOLOGY LLC [US/US]; 800 N. Lind-  
bergh Boulevard, St. Louis, MO 63167 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): DEHESH, Katay-  
oon [US/US]; 521 Crownpointe Circle, Vacaville, CA  
95687 (US). KNAUF, Vic, C. [US/US]; 11835 NE Yeomalt  
Point Drive, Bainbridge Island, WA 98110 (US).

(74) Agents: MARSH, David, R. et al.; Arnold & Porter, Attn:  
IP Docketing Dept., Room 1126B, 555 Twelfth Street,  
N.W., Washington, DC 20004-1206 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,  
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,  
MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC,  
SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG,  
US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,  
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,  
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,  
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Published:**

— without international search report and to be republished  
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: THIOESTERASE-RELATED NUCLEIC ACID SEQUENCES AND METHODS OF USE FOR THE PRODUCTION  
OF PLANTS WITH MODIFIED FATTY ACID COMPOSITION

(57) Abstract: The present invention is directed to nucleic acid molecules and nucleic acid constructs, and other agents associated  
with fatty acid synthesis, particularly the ratios of saturated and unsaturated fats. Moreover, the present invention is directed to plants  
incorporating such agents where the plants exhibit altered ratios of saturated and unsaturated fats. In particular, the present invention  
is directed to plants incorporating such agents where the plants exhibit altered levels of saturated and unsaturated fatty acids.

WO 2004/000871 A2

**THIOESTERASE-RELATED NUCLEIC ACID SEQUENCES  
AND METHODS OF USE FOR THE PRODUCTION OF PLANTS  
WITH MODIFIED FATTY ACID COMPOSITION**

5

**FIELD OF THE INVENTION**

The present invention is directed to nucleic acid molecules and nucleic acid constructs, and other agents associated with fatty acid synthesis. Moreover, the present invention is directed to plants incorporating such agents where the plants exhibit altered ratios of saturated and unsaturated fats. In particular, the present invention is directed to plants incorporating such agents where the plants exhibit altered ratios of saturated to unsaturated fatty acids.

**BACKGROUND**

Plant oils are used in a variety of applications. Novel vegetable oil compositions and improved approaches to obtain oil compositions, from biosynthetic or natural plant sources, are needed. Depending upon the intended oil use, various different fatty acid compositions are desired. Plants, especially plant species which synthesize large amounts of oils in plant seeds, are an important source of oils both for edible and industrial uses.

With the exception of coconut endosperm and palm kernel oils, which contain high amounts of laurate (C12:0), the common edible oils all basically consist of palmitate (16:0), stearate (18:0), oleate (18:1), linoleate (18:2), and linolenate (18:3). Many oilseed species have highly elevated levels of saturated fatty acids. Coconut oil contains more than 90% saturated fatty acids, predominantly laurate (12:0) and other medium chain fatty acids ranging from C6 to C16. Other highly saturated oils include oils with high palmitate (16:0) and stearate (18:0) levels (up to approximately 60% of acyl chains). These oils include those derived from cocoa butter (25% palmitate; 34% stearate) and oil palm mesocarp (45% palmitate; 15% stearate). Soybean oil typically contains about 16-20% saturated fatty acids: 13-16% palmitate and 3-4% stearate. Voelker *et al.*, 52 *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 335-61 (2001).

For many oil applications, saturated fatty acid levels are preferably less than 6% by weight, and more preferably about 2-3% by weight. Saturated fatty acids have undesirable

high melting points and cloud at low temperatures. Products created from oils containing low saturated fatty acid levels may be preferred by consumers and the food industry because they are perceived to be healthier and/or may be labeled as "saturated fat free" or "trans fat free" products in accordance with FDA guidelines. Oils with low saturated fatty acid levels have improved cold flow properties, which are important in biodiesel and lubricant applications, and do not cloud at low temperatures, thereby reducing or eliminating the need to winterize the oil for food applications such as salad oils.

Higher plants synthesize fatty acids in the plastids via the fatty acid synthetase (FAS) pathway. In developing oilseeds, most fatty acids are attached to glycerol backbones to form triglycerides, for storage as a source of energy.

$\beta$ -ketoacyl-ACP synthase I catalyzes elongation up to palmitoyl-ACP (C16:0), whereas  $\beta$ -ketoacyl-ACP synthase II catalyzes the final elongation to stearyl-ACP (C18:0). Common plant unsaturated fatty acids, such as oleic, linoleic and linolenic acids found in storage triglycerides, originate from the desaturation of stearyl-ACP to form oleoyl-ACP (C18:1) in a reaction catalyzed by a soluble plastid delta-9 desaturase (also often referred to as "stearyl-ACP desaturase"). Additional desaturation is effected sequentially by the actions of membrane bound delta-12 desaturase and delta-15 desaturase. These "desaturases" thus create polyunsaturated fatty acids.

Specific thioesterases can terminate fatty acid chain elongation by hydrolyzing the newly produced acyl-ACPs into free fatty acids and ACP. Subsequently, the free fatty acids are converted to fatty acyl-CoAs in the plastid envelope and exported to the cytoplasm, where they may be incorporated into the endoplasmic reticulum (ER) lipid biosynthesis pathway (Kennedy pathway), which is responsible for the formation of phospholipids, triglycerides, and other neutral lipids. Plant acyl-ACP thioesterases are of biochemical interest because of their roles in fatty acid synthesis and their utility in bioengineering of plant oil seeds. The thioesterases have an important role in determining chain length during de novo fatty acid biosynthesis in plants, and thus these enzymes are useful in the provision of various modifications of fatty acyl compositions, particular with respect to the relative proportions of various fatty acyl groups that are present in seed storage oils.

Plant thioesterases can be classified into two gene families based on sequence homology and substrate preference. The first gene family, FATA, includes long chain acyl-ACP thioesterases having activity primarily on oleoyl-ACP (18:1-ACP). Oleoyl-ACP is the immediate precursor of most fatty acids found in phospholipids and triglycerides synthesized

by the ER lipid biosynthetic pathway. A second class of plant thioesterases, FATB, includes enzymes that, in most plants, utilize palmitoyl-ACP (16:0-ACP), stearoyl (18:0-ACP), and oleoyl-ACP (18:1-ACP). FATB enzymes have been isolated from California bay (*Umbellularia californica*) (U.S. Patent No. 5,955,329; U.S. Patent No. 5,723,761), elm (U.S. Patent No. 5,723,761), *Cuphea hookeriana* (U.S. Patent No. 5,723,761), *Cuphea palustris* (U.S. Patent No. 5,955,329), *Cuphea lanceolata*, nutmeg, *Arabidopsis thaliana*, mango (U.S. Patent No. 5,723,761), leek (U.S. Patent No. 5,723,761), camphor (*Cinnamomum camphora*) (U.S. Patent No. 5,955,329), canola (U.S. Patent No. 5,955,650), and maize (U.S. Patent No. 6,331,664).

Obtaining nucleic acid sequences capable of producing a phenotypic result in FAS, desaturation and/or incorporation of fatty acids into a glycerol backbone to produce an oil is subject to various obstacles including but not limited to the identification of metabolic factors of interest, choice and characterization of an enzyme source with useful kinetic properties, purification of the protein of interest to a level which will allow for its amino acid sequencing, utilizing amino acid sequence data to obtain a nucleic acid sequence capable of use as a probe to retrieve the desired DNA sequence, and the preparation of gene constructs, transformation and analysis of the resulting plants.

Thus, additional nucleic acid targets and methods for modifying fatty acid compositions are needed. In particular, constructs and methods to produce a variety of ranges of different fatty acid compositions are needed.

## SUMMARY OF THE INVENTION

The present invention provides a substantially purified nucleic acid molecule comprising a nucleic acid sequence with at least 70% sequence identity to SEQ ID NO: 2 or its complement. Also provided by the present invention is a substantially purified nucleic acid molecule comprising a nucleic acid sequence with at least 70% sequence identity to SEQ ID NO: 3 or its complement. Also provided by the present invention is a substantially purified nucleic acid molecule comprising a nucleic acid sequence with at least 70% sequence identity to SEQ ID NO: 4 or its complement. Also provided by the present invention is a substantially purified nucleic acid molecule comprising a nucleic acid sequence with at least 70% sequence identity to SEQ ID NO: 5 or its complement. Also provided by the present invention is a substantially purified nucleic acid molecule comprising a nucleic

acid sequence with at least 70% sequence identity to SEQ ID NO: 6 or its complement. Also provided by the present invention is a substantially purified nucleic acid molecule comprising a nucleic acid sequence with at least 70% sequence identity to SEQ ID NO: 7 or its complement. Also provided by the present invention is a substantially purified nucleic acid molecule comprising a nucleic acid sequence with at least 70% sequence identity to SEQ ID NO: 8 or its complement. Further provided by the present invention are a nucleic acid molecule comprising at least 15 consecutive nucleotides of a nucleic acid molecule having the sequence of SEQ ID NO: 2; and a nucleic acid molecule comprising at least 15 consecutive nucleotides of a nucleic acid molecule having the sequence of SEQ ID NO: 3; and a nucleic acid molecule comprising at least 15 consecutive nucleotides of a nucleic acid molecule having the sequence of SEQ ID NO: 4; and a nucleic acid molecule comprising at least 15 consecutive nucleotides of a nucleic acid molecule having the sequence of SEQ ID NO: 5; and a nucleic acid molecule comprising at least 15 consecutive nucleotides of a nucleic acid molecule having the sequence of SEQ ID NO: 6; and a nucleic acid molecule comprising at least 15 consecutive nucleotides of a nucleic acid molecule having the sequence of SEQ ID NO: 7; and a nucleic acid molecule comprising at least 15 consecutive nucleotides of a nucleic acid molecule having the sequence of SEQ ID NO: 8.

Also provided by the present invention is a recombinant nucleic acid molecule comprising as operably linked components: (A) a promoter that functions in a plant cell to cause production of an mRNA molecule; and (B) a nucleic acid sequence that has at least 85% identity to a nucleic acid sequence selected from the group consisting of SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, complements thereof, and fragments of either.

Also provided by the present invention is an intron obtained from a genomic polynucleotide sequence wherein the genomic polynucleotide sequence is selected from the group consisting of: (a) a genomic polynucleotide sequence having at least 70% identity to coding regions of SEQ ID NO: 1 over the entire length of SEQ ID NO: 1; (b) a genomic polynucleotide sequence having at least 80% identity to coding regions of SEQ ID NO: 1 over the entire length of SEQ ID NO: 1; (c) a genomic polynucleotide sequence having at least 90% identity to coding regions of SEQ ID NO: 1 over the entire length of SEQ ID NO: 1; and (d) a genomic polynucleotide sequence having at least 95% identity to coding regions of SEQ ID NO: 1 over the entire length of SEQ ID NO: 1.

Also provided by the present invention is an intron obtained from a genomic polynucleotide sequence wherein the genomic polynucleotide sequence is selected from the group consisting of: (a) a genomic polynucleotide sequence having at least 70% identity to coding regions of SEQ ID NO: 10 over the entire length of SEQ ID NO: 10; (b) a genomic polynucleotide sequence having at least 80% identity to coding regions of SEQ ID NO: 10 over the entire length of SEQ ID NO: 10; (c) a genomic polynucleotide sequence having at least 90% identity to coding regions of SEQ ID NO: 10 over the entire length of SEQ ID NO: 10; and (d) a genomic polynucleotide sequence having at least 95% identity to coding regions of SEQ ID NO: 10 over the entire length of SEQ ID NO: 10.

Also provided by the present invention are transformed plant cells and plants comprising a recombinant nucleic acid molecule, the recombinant nucleic acid molecule comprising as operably linked components: (A) a promoter that functions in a plant cell to cause production of an mRNA molecule; and (B) a nucleic acid sequence that has at least 85% identity to a nucleic acid sequence selected from the group consisting of SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, complements thereof, and fragments of either.

The present invention also provides a transformed soybean plant having a recombinant nucleic acid molecule that comprises a promoter operably linked to a nucleic acid sequence that has at least 85% identity to a nucleic acid sequence selected from the group consisting of SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, complements thereof, and fragments of either.

Further provided by the present invention is a transformed soybean plant having a nucleic acid molecule that comprises (a) a first promoter operably linked to a first nucleic acid molecule having a first nucleic acid sequence that has 85% or greater identity to a nucleic acid sequence selected from the group consisting of SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, complements thereof, and fragments of either, and (b) a second nucleic acid molecule having a second nucleic acid sequence that encodes an enzyme selected from the group consisting of beta-ketoacyl-ACP synthase I, beta-ketoacyl-ACP synthase IV, and delta-9 desaturase.

The present invention also provides seed derived from a transformed plant which comprises a recombinant nucleic acid molecule, the recombinant nucleic acid molecule comprising as operably linked components: (A) a promoter that functions in a plant to cause production of an mRNA molecule; and (B) a nucleic acid sequence that has at least 85%

identity to a nucleic acid sequence selected from the group consisting of SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, complements thereof, and fragments of either.

Also provided by the present invention is oil derived from seed of a transformed  
5 plant, where the transformed plant comprises a recombinant nucleic acid molecule which comprises as operably linked components: (A) a promoter that functions in a plant to cause production of an mRNA molecule; and (B) a nucleic acid sequence that has at least 85% identity to a nucleic acid sequence selected from the group consisting of SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8,  
10 complements thereof, and fragments of either, where the oil exhibits a reduced saturated fatty acid content relative to oil derived from seed of a plant with a similar genetic background but lacking the recombinant nucleic acid molecule.

The present invention also provides a method of producing a transformed plant having seed with a reduced saturated fatty acid content comprising: (A) transforming a plant with a  
15 nucleic acid molecule to produce a transformed plant, where the nucleic acid molecule comprises a nucleic acid sequence that has 85% or greater identity to an intron of a plant thioesterase gene; and (B) growing the transformed plant, where the plant produces seed with a reduced saturated fatty acid content relative to a plant having a similar genetic background but lacking the nucleic acid molecule.

Further provided by the present invention is a method of producing a plant having a seed with reduced palmitic and stearic acid levels comprising: transforming a plant with a nucleic acid molecule that comprises (a) a first promoter operably linked to a first nucleic acid molecule having a first nucleic acid sequence that has 85% or greater identity to a nucleic acid sequence selected from the group consisting of SEQ ID NO: 2, SEQ ID NO: 3,  
25 SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, complements thereof, and fragments of either, and (b) a second nucleic acid molecule having a second nucleic acid sequence that encodes an enzyme selected from the group consisting of beta-ketoacyl-ACP synthase I, beta-ketoacyl-ACP synthase IV, and delta-9 desaturase; and growing the plant, where the plant produces seed with reduced palmitic and stearic acid  
30 levels relative to a plant having a similar genetic background but lacking the nucleic acid molecule.

The present invention also provides a method of producing a plant having a seed with a modified oil composition comprising: transforming a plant with a nucleic acid molecule

that comprises, as operably linked components, a first promoter and a first nucleic acid molecule having a first nucleic acid sequence that has 85% or greater identity to a nucleic acid sequence selected from the group consisting of SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, complements thereof, and fragments of either; and, growing the plant, where the plant produces seed with a modified oil composition relative to a plant having a similar genetic background but lacking the nucleic acid molecule.

The present invention further provides a method of modifying the lipid composition in a plant cell comprising: transforming a plant cell with a recombinant DNA construct having a DNA sequence selected from the group consisting of SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, complements thereof, and fragments of either, and growing the cell under conditions where transcription of the DNA sequence is initiated, where the lipid composition is modified.

Also provided by the present invention is a method of modifying the lipid composition in a host cell comprising: transforming a host cell with a DNA construct comprising as operably associated components in the 5' to 3' direction of transcription, a transcriptional initiation region functional in the host cell, a DNA sequence selected from the group consisting of SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, complements thereof, and fragments of either, and a transcription termination sequence, and growing the cell under conditions where transcription of the DNA sequence is initiated, where the lipid composition is modified.

Further provided by the present invention is a method of altering the expression of a *FATB* gene in a seed comprising: (a) introducing into a plant cell a first DNA sequence capable of expressing a first RNA that exhibits at least 90% identity to a transcribed intron of the *FATB* gene, and a second DNA sequence capable of expressing a second RNA capable of forming a dsRNA with the first RNA; and (b) expressing the first RNA and the second RNA in a seed.

Also provided by the present invention are methods of altering the expression of a *FATB* gene in a seed comprising: (a) introducing into a plant cell a first DNA sequence capable of expressing an RNA that exhibits at least 90% identity to a transcribed intron of the *FATB* gene and a second DNA sequence capable of expressing a second RNA that exhibits at least 90% identity to a transcribed intron of the *FATB* gene; and (b) expressing said the first RNA and the second RNA in a seed.

## BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a schematic of construct pCGN3892.

Figure 2 is a schematic of construct pMON70674.

Figure 3 is a schematic of construct pMON41164.

5 Figure 4 is a schematic of construct pMON70678.

Figure 5 is a schematic of construct pMON70675.

Figure 6 is a schematic of construct pMON70680.

Figure 7 is a schematic of construct pMON70656.

Figure 8 is a schematic of construct pMON70681.

## 10 DETAILED DESCRIPTION OF THE INVENTION

Description of the Nucleic Acid Sequences

SEQ ID NO: 1 sets forth a nucleic acid sequence of a soybean *FATB* genomic clone.

SEQ ID NO: 2 sets forth a nucleic acid sequence of a soybean *FATB* intron I.

SEQ ID NO: 3 sets forth a nucleic acid sequence of a soybean *FATB* intron II.

15 SEQ ID NO: 4 sets forth a nucleic acid sequence of a soybean *FATB* intron III.

SEQ ID NO: 5 sets forth a nucleic acid sequence of a soybean *FATB* intron IV.

SEQ ID NO: 6 sets forth a nucleic acid sequence of a soybean *FATB* intron V.

SEQ ID NO: 7 sets forth a nucleic acid sequence of a soybean *FATB* intron VI.

SEQ ID NO: 8 sets forth a nucleic acid sequence of a soybean *FATB* intron VII.

20 SEQ ID NO: 9 sets forth an amino acid sequence of a soybean *FATB* enzyme.

SEQ ID NO: 10 sets forth a nucleic acid sequence of a soybean *FATB* partial genomic clone.

SEQ ID NOs: 11-18 set forth nucleic acid sequences of oligonucleotide primers.

25 SEQ ID NO: 19 sets forth a nucleic acid sequence of a PCR product containing soybean *FATB* intron II.

SEQ ID NO: 20 sets forth a nucleic acid sequence of a soybean *FATB* cDNA.

Definitions

30 As used herein, the term "gene" is used to refer to the nucleic acid sequence that encompasses the 5' promoter region associated with the expression of the gene product, any

intron and exon regions and 3' untranslated regions associated with the expression of the gene product.

As used herein, the term "ACP" is used to refer to an acyl carrier protein moiety. The term "fatty acid", as used herein, refers to free fatty acids and acyl-fatty acid groups.

5 As used herein, a "*FATB*" or "palmitoyl-ACP thioesterase" gene is a gene that encodes an enzyme (FATB) capable of catalyzing the hydrolytic cleavage of the carbon-sulfur thioester bond in the panthothene prosthetic group of palmitoyl-ACP as its preferred reaction. Hydrolysis of other fatty acid-ACP thioesters may also be catalyzed by this enzyme.

10 When referring to proteins and nucleic acids herein, the use of plain capitals, *e.g.*, "*FATB*", indicates a reference to an enzyme, protein, polypeptide, or peptide, and the use of italicized capitals, *e.g.*, "*FATB*", is used to refer to nucleic acids, including without limitation genes, cDNAs, and mRNAs.

As used herein, a "beta-ketoacyl-ACP synthase I" or "*KAS I*" gene is a gene that  
15 encodes an enzyme (KAS I) capable of catalyzing the elongation of a fatty acyl moiety up to palmitoyl-ACP (C16:0). Exemplary KAS I genes and enzymes are described in U.S. Patent No. 5,475,099 and PCT Publication WO 94/10189.

As used herein, a "beta-ketoacyl-ACP synthase IV" or "*KAS IV*" gene is a gene that  
encodes an enzyme (KAS IV) capable of catalyzing the condensation of medium chain acyl-  
20 ACPs. Exemplary KAS IV genes and enzymes are described in PCT Publication WO 98/46776.

As used herein, a "delta-9 desaturase" or "stearoyl-ACP desaturase" or "omega-9  
desaturase" gene is a gene that encodes an enzyme capable of catalyzing the insertion of a  
double bond into a fatty acyl moiety at the ninth position counted from the carboxyl terminus.  
25 Exemplary delta-9 desaturase genes and enzymes are described in U.S. Patent No. 5,723,595.

As used herein, a "mid-oleic soybean seed" is a seed having between 50% and 75%  
oleic acid present in the oil composition of the seed.

As used herein, a "high oleic soybean seed" is a seed with oil having greater than 75%  
oleic acid present in the oil composition of the seed.

30 As used herein, a "low saturate" oil composition contains between 3.4 and 7 percent  
saturated fatty acids.

As used herein, a "zero saturate" oil composition contains less than 3.4 percent  
saturated fatty acids.

As used herein, a cell or organism can have a family of more than one gene encoding a particular enzyme, e.g., a plant can have a family of more than one *FATB* gene (*i.e.*, genes which encode an enzyme with the specified activity present at different locations within the genome of the plants). As used herein, a "*FATB* gene family member" is any *FATB* gene found within the genetic material of the plant. In one embodiment, a gene family can be additionally classified by the similarity of the nucleic acid sequences. In a preferred aspect of this embodiment, a gene family member exhibits at least 60%, more preferably at least 70%, more preferably at least 80% nucleic acid sequence identity in the coding sequence portion of the gene.

The term "non-coding" refers to sequences of nucleic acid molecules that do not encode part or all of an expressed protein. Non-coding sequences include but are not limited to introns, promoter regions, 3' untranslated regions, and 5' untranslated regions.

The term "intron" as used herein refers to the normal sense of the term as meaning a segment of nucleic acid molecules, usually DNA, that does not encode part of or all of an expressed protein, and which, in endogenous conditions, is transcribed into RNA molecules, but which is spliced out of the endogenous RNA before the RNA is translated into a protein.

The term "exon" as used herein refers to the normal sense of the term as meaning a segment of nucleic acid molecules, usually DNA, that encodes part of or all of an expressed protein.

As used herein, a promoter that is "operably linked" to one or more nucleic acid sequences is capable of driving expression of one or more nucleic acid sequences, including multiple coding or non-coding nucleic acid sequences arranged in a polycistronic configuration.

A "polycistronic gene" or "polycistronic mRNA" is any gene or mRNA that contains transcribed nucleic acid sequences which correspond to nucleic acid sequences of more than one gene targeted for expression. It is understood that such polycistronic genes or mRNAs may contain sequences that correspond to introns, 5'UTRs, 3'UTRs, or combinations thereof, and that a recombinant polycistronic gene or mRNA might, for example without limitation, contain sequences that correspond to one or more UTRs from one gene and one or more introns from a second gene.

As used herein, the term complement of a nucleic acid sequence refers to the complement of the sequence along its complete length.

As used herein, any range set forth is inclusive of the end points of the range unless otherwise stated.

### Agents

The agents of the invention will preferably be "biologically active" with respect to either a structural attribute, such as the capacity of a nucleic acid molecule to hybridize to another nucleic acid molecule, or the ability of a protein to be bound by an antibody (or to compete with another molecule for such binding). Alternatively, such an attribute may be catalytic and thus involve the capacity of the agent to mediate a chemical reaction or response. The agents will preferably be "substantially purified." The term "substantially purified," as used herein, refers to a molecule separated from substantially all other molecules normally associated with it in its native environmental conditions. More preferably a substantially purified molecule is the predominant species present in a preparation. A substantially purified molecule may be greater than 60% free, greater than 75% free, preferably greater than 90% free, and most preferably greater than 95% free from the other molecules (exclusive of solvent) present in the natural mixture. The term "substantially purified" is not intended to encompass molecules present in their native environmental conditions.

The agents of the invention may also be recombinant. As used herein, the term "recombinant" means any agent (*e.g.*, including but limited to DNA, peptide), that is, or results, however indirectly, from human manipulation of a nucleic acid molecule.

It is understood that the agents of the invention may be labeled with reagents that facilitate detection of the agent (*e.g.*, fluorescent labels, Prober *et al.*, *Science* 238:336-340 (1987); Albarella *et al.*, EP 144914; chemical labels, Sheldon *et al.*, U.S. Patent 4,582,789; Albarella *et al.*, U.S. Patent 4,563,417; modified bases, Miyoshi *et al.*, EP 119448).

### Nucleic Acid Molecules

Agents of the invention include nucleic acid molecules. In an aspect of the present invention, the nucleic acid molecule comprises a nucleic acid sequence, which when introduced into a cell or organism, is capable of selectively reducing the level of a FATB protein and/or *FATB* transcript.

In a preferred aspect of the invention, the nucleic acid sequences are intron sequences or other non-coding sequences of a *FATB* gene, which when introduced into a cell or organism are capable of selectively reducing the level of an endogenous *FATB* protein and/or endogenous *FATB* transcript, thereby resulting in a modification of the fatty acid biosynthetic

pathway and a consequent decrease in the levels of saturated fatty acids in the cell or organism. Non-coding sequences of a *FATB* gene may also be used in combination with nucleic acid sequences coding for enzymes such as beta-ketoacyl-ACP synthase I, beta-ketoacyl-ACP synthase IV, and delta-9 desaturase, which further modifies the fatty acid biosynthetic pathway and further decreases the level of saturated fatty acids in the cell or organism. Non-coding sequences of a *FATB* gene may also be used in combination with nucleic acid sequences that down-regulate other enzymes, for example a cDNA that is capable of sense suppression of a delta-12 desaturase gene, which further modifies the fatty acid biosynthetic pathway and further decreases the level of saturated fatty acids in the cell or organism.

In a preferred aspect, the capability of a nucleic acid molecule to selectively reduce the level of a protein and/or transcript is carried out by a comparison of levels of mRNA transcripts. In another preferred aspect of the present invention, the nucleic acid molecule of the invention comprises a nucleic acid sequence selected from the group consisting of SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, complements thereof, and fragments of either.

In one aspect of the present invention the nucleic acids of the present invention are said to be introduced nucleic acid molecules. A nucleic acid molecule is said to be "introduced" if it is inserted into a cell or organism as a result of human manipulation, no matter how indirect. Examples of introduced nucleic acid molecules include, but are not limited to, nucleic acids that have been introduced into cells via transformation, transfection, injection, and projection, and those that have been introduced into an organism via methods including, but not limited to, conjugation, endocytosis, and phagocytosis. The cell or organism can be, or can be derived from, without limitation, a plant, plant cell, algae cell, algae, fungal cell, fungus, or bacterial cell.

As used herein, "a selective reduction" of an agent such as a protein, fatty acid, or mRNA is relative to a cell or organism lacking a nucleic acid molecule capable of selectively reducing the agent. In a preferred aspect, the level of the agent is selectively reduced by at least 50%, preferably at least more than 75%, and even more preferably by at least more than 90% or 95%.

As used herein, "a partial reduction" of the level of an agent such as a protein, fatty acid, or mRNA means that the level is reduced at least 25% relative to a cell or organism lacking a nucleic acid molecule capable of reducing the agent.

As used herein, "a substantial reduction" of the level of an agent such as a protein, fatty acid, or mRNA means that the level is reduced at least 75% relative to a cell or organism lacking a nucleic acid molecule capable of reducing the agent.

As used herein, "an effective elimination" of an agent such as a protein, fatty acid, or mRNA means that the level of the agent is reduced at least 95% relative to a cell or organism lacking a nucleic acid molecule capable of reducing the agent.

When levels of an agent are compared, such a comparison is preferably carried out between organisms with a similar genetic background. In a preferred aspect, a similar genetic background is a background where the organisms being compared share 50% or greater of their nuclear genetic material. In a more preferred aspect a similar genetic background is a background where the organisms being compared share 75% or greater, even more preferably 90% or greater of their nuclear genetic material. In another even more preferable aspect, a similar genetic background is a background where the organisms being compared are plants, and the plants are isogenic except for any genetic material originally introduced using plant transformation techniques.

In an embodiment of the present invention, a nucleic acid molecule, when introduced into a cell or organism, is capable of selectively reducing the level of a protein, fatty acid, and/or transcript. In a preferred aspect, the capability of a nucleic acid molecule to selectively reduce the level of a protein, fatty acid, and/or transcript is determined relative to a cell or organism lacking a nucleic acid molecule capable of selectively reducing the protein, fatty acid, and/or transcript. As used herein, mRNA transcripts include processed and non-processed mRNA transcripts, and a "*FATB* transcript" refers to any transcript encoded by a *FATB* gene.

In another embodiment, a nucleic acid molecule, when introduced into a cell or organism, is capable of at least partially reducing the level of a *FATB* protein and/or *FATB* transcript. In a different embodiment, a nucleic acid molecule, when introduced into a cell or organism, is capable of at least substantially reducing the level of a *FATB* protein and/or *FATB* transcript. In a further embodiment, a nucleic acid molecule, when introduced into a cell or organism, is capable of effectively eliminating the level of a *FATB* protein and/or *FATB* transcript.

In a different embodiment, a nucleic acid molecule, when introduced into a cell or organism, is capable of selectively reducing the level of a *FATB* protein and/or *FATB* transcript while overexpressing the level of a different protein and/or transcript. Preferably,

the different protein is selected from the group consisting of beta-ketoacyl-ACP synthase I, beta-ketoacyl-ACP synthase IV, and delta-9 desaturase, and the different transcript encodes an enzyme selected from the group consisting of beta-ketoacyl-ACP synthase I, beta-ketoacyl-ACP synthase IV, and delta-9 desaturase.

- 5 In a further embodiment, a nucleic acid molecule, when introduced into a cell or organism, is capable of at least partially reducing the level of a FATB protein and/or *FATB* transcript while overexpressing the level of a different protein and/or transcript. Preferably, the different protein is selected from the group consisting of beta-ketoacyl-ACP synthase I, beta-ketoacyl-ACP synthase IV, and delta-9 desaturase, and the different transcript encodes
- 10 an enzyme selected from the group consisting of beta-ketoacyl-ACP synthase I, beta-ketoacyl-ACP synthase IV, and delta-9 desaturase. In a different embodiment, a nucleic acid molecule, when introduced into a cell or organism, is capable of at least substantially reducing the level of a FATB protein and/or *FATB* transcript while overexpressing the level of a different protein and/or transcript. In a further embodiment, a nucleic acid molecule,
- 15 when introduced into a cell or organism, is capable of effectively eliminating the level of a FATB protein and/or *FATB* transcript while overexpressing the level of a different protein and/or transcript.

- Further preferred embodiments of the invention are nucleic acid molecules that are at least 50%, 60%, or 70% identical over their entire length to a nucleic acid molecule of the
- 20 invention, and nucleic acid molecules that are complementary to such nucleic acid molecules. More preferable are nucleic acid molecules that comprise a region that is at least 80% or 85% identical over its entire length to a nucleic acid molecule of the invention and nucleic acid molecules that are complementary thereto. In this regard, nucleic acid molecules at least 90% identical over their entire length are particularly preferred, those at least 95% identical are
- 25 especially preferred. Further, those with at least 97% identity are highly preferred and those with at least 98% and 99% identity are particularly highly preferred, with those at least 99% being the most highly preferred.

- The invention also provides a nucleic acid molecule comprising a nucleic acid molecule sequence obtainable by screening an appropriate library containing the complete
- 30 gene for a nucleic acid molecule sequence set forth in the Sequence Listing under stringent hybridization conditions with a probe having the sequence of said nucleic acid molecule sequence or a fragment thereof; and isolating said nucleic acid molecule sequence.

Fragments useful for obtaining such a nucleic acid molecule include, for example, probes and primers as described herein.

Nucleic acid molecules of the invention can be used as a hybridization probe for RNA, cDNA, or genomic DNA to isolate full length cDNAs or genomic clones and to isolate  
5 cDNA or genomic clones of other genes that have a high sequence similarity to a nucleic acid molecule set forth in the Sequence Listing.

The nucleic acid molecules of the present invention can be readily obtained by using the herein described nucleic acid molecules or fragments thereof to screen cDNA or genomic libraries obtained from plant species or other appropriate organisms. These methods are  
10 known to those of skill in the art, as are methods for forming such libraries. In one embodiment, such sequences are obtained by incubating nucleic acid molecules of the present invention with members of genomic libraries and recovering clones that hybridize to such nucleic acid molecules thereof. In a second embodiment, methods of chromosome walking or inverse PCR may be used to obtain such sequences. In a third embodiment, the sequence  
15 of a nucleic acid molecule of the present invention may be used to screen a library or database, using bioinformatics techniques known in the art. *See, e.g., Bioinformatics*, Baxevanis & Ouellette, eds., Wiley-Interscience (1998).

Any of a variety of methods may be used to obtain one or more of the nucleic acid molecules of the present invention. Automated nucleic acid synthesizers may be employed  
20 for this purpose, and to make a nucleic acid molecule that has a sequence also found in a cell or organism. In lieu of such synthesis, the disclosed nucleic acid molecules may be used to define a pair of primers that can be used with the polymerase chain reaction to amplify and obtain any desired nucleic acid molecule or fragment.

"Identity," as is well understood in the art, is a relationship between two or more  
25 polypeptide sequences or two or more nucleic acid molecule sequences, as determined by comparing the sequences. In the art, "identity" also means the degree of sequence relatedness between polypeptide or nucleic acid molecule sequences, as determined by the match between strings of such sequences. "Identity" can be readily calculated by known methods including, but not limited to, those described in *Computational Molecular Biology*, Lesk,  
30 A.M., ed., Oxford University Press, New York (1988); *Biocomputing: Informatics and Genome Projects*, Smith, D.W., ed., Academic Press, New York (1993); *Computer Analysis of Sequence Data, Part I*, Griffin, A.M. and Griffin, H.G., eds., Humana Press, New Jersey (1994); *Sequence Analysis in Molecular Biology*, von Heinje, G., Academic Press (1987);

*Sequence Analysis Primer*, Gribskov, M. and Devereux, J., eds., Stockton Press, New York (1991); and Carillo, H., and Lipman, D., *SIAM J. Applied Math*, 48:1073 (1988). Methods to determine identity are designed to give the largest match between the sequences tested.

Moreover, methods to determine identity are codified in publicly available programs.

- 5 Computer programs which can be used to determine identity between two sequences include, but are not limited to, GCG (Devereux, J., *et al.*, *Nucleic Acids Research* 12(1):387 (1984); suite of five BLAST programs, three designed for nucleotide sequences queries (BLASTN, BLASTX, and TBLASTX) and two designed for protein sequence queries (BLASTP and TBLASTN) (Coulson, *Trends in Biotechnology*, 12:76-80 (1994); Birren *et al.*, *Genome*  
10 *Analysis*, 1:543-559 (1997)). The BLASTX program is publicly available from NCBI and other sources (*BLAST Manual*, Altschul, S., *et al.*, NCBI NLM NIH, Bethesda, MD 20894; Altschul, S., *et al.*, *J. Mol. Biol.*, 215:403-410 (1990)). The well-known Smith Waterman algorithm can also be used to determine identity.

Parameters for polypeptide sequence comparison typically include the following:

- 15 Algorithm: Needleman and Wunsch, *J. Mol. Biol.*, 48:443-453 (1970)  
Comparison matrix: BLOSSUM62 from Hentikoff and Hentikoff, *Proc. Natl. Acad. Sci. USA*, 89:10915-10919 (1992)

Gap Penalty: 12

Gap Length Penalty: 4

- 20 A program which can be used with these parameters is publicly available as the "gap" program from Genetics Computer Group, Madison, Wisconsin. The above parameters along with no penalty for end gap are the default parameters for peptide comparisons.

Parameters for nucleic acid molecule sequence comparison include the following:

- Algorithm: Needleman and Wunsch, *J. Mol. Bio.*, 48:443-453 (1970)

- 25 Comparison matrix: matches - +10; mismatches = 0

Gap Penalty: 50

Gap Length Penalty: 3

- As used herein, "% identity" is determined using the above parameters as the default parameters for nucleic acid molecule sequence comparisons and the "gap" program from  
30 GCG, version 10.2.

The invention further relates to nucleic acid molecules that hybridize to nucleic acid molecules of the present invention. In particular, the invention relates to nucleic acid molecules that hybridize under stringent conditions to the above-described nucleic acid

molecules. As used herein, the terms "stringent conditions" and "stringent hybridization conditions" mean that hybridization will generally occur if there is at least 95% and preferably at least 97% identity between the sequences. An example of stringent hybridization conditions is overnight incubation at 42°C in a solution comprising 50%

5 formamide, 5x SSC (150 mM NaCl, 15 mM trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 micrograms/ milliliter denatured, sheared salmon sperm DNA, followed by washing the hybridization support in 0.1x SSC at approximately 65°C. Other hybridization and wash conditions are well known and are exemplified in Sambrook *et al.*, Molecular Cloning: A Laboratory Manual, Second Edition, 10 Cold Spring Harbor, NY (1989), particularly Chapter 11.

In embodiments where nucleic acid sequences which when expressed are capable of selectively reducing the level of a *FATB* protein and/or *FATB* transcript, preferred nucleic acid sequences are selected from the groups consisting of (1) nucleic acid sequences with at least 50%, 60%, 70%, 80%, 90%, 95%, 97%, 98%, 99% or 100% sequence identity over the 15 entire length of the nucleic acid molecule with a nucleotide sequence selected from the group consisting of SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, complements thereof, and fragments of either; (2) nucleic acid molecules which contain sequences that are also found in a soybean *FATB* gene intron; and (3) nucleic acid molecules that exhibit sequences with at least 50%, 60%, 70%, 80%, 20 85%, 90%, 95%, 97%, 98%, 99% or 100% sequence identity over the entire length of the nucleic acid molecule with a nucleic acid molecule of (2).

One subset of the nucleic acid molecules of the invention includes fragment nucleic acid molecules. Fragment nucleic acid molecules may consist of significant portion(s) of, or indeed most of, the nucleic acid molecules of the invention, such as those specifically 25 disclosed. Alternatively, the fragments may comprise smaller oligonucleotides (having from about 15 to about 400 contiguous nucleotide residues and more preferably, about 15 to about 30 contiguous nucleotide residues, or about 50 to about 100 contiguous nucleotide residues, or about 100 to about 200 contiguous nucleotide residues, or about 200 to about 400 contiguous nucleotide residues, or about 275 to about 350 contiguous nucleotide residues). 30 More preferably, the fragments may comprise small oligonucleotides having from about 15 to about 45 contiguous nucleotide residues, about 20 to about 45 contiguous nucleotide residues, about 15 to about 30 contiguous nucleotide residues, about 21 to about 30 contiguous nucleotide residues, about 21 to about 25 contiguous nucleotide residues, about 21 to about

25 contiguous nucleotide residues, about 19 to about 25 contiguous nucleotide residues, or about 21 contiguous nucleotides.

In another aspect, a fragment nucleic acid molecule has a nucleic acid sequence that is at least 15, 25, 50, or 100 contiguous nucleotides of a nucleic acid molecule of the present invention. In a preferred embodiment, the nucleic acid molecule has a nucleic acid sequence  
5 that is at least 15, 25, 50, or 100 contiguous nucleotides of a nucleic acid molecule having a nucleic acid sequence selected from the group consisting of SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, and complements thereof.

10 A fragment of one or more of the nucleic acid molecules of the present invention may be a probe and specifically a PCR probe. A PCR probe is a nucleic acid molecule capable of initiating a polymerase activity while in a double-stranded structure with another nucleic acid molecule. Various methods for determining the structure of PCR probes and PCR techniques exist in the art. Computer generated searches using programs such as Primer3 ([www-genome.wi.mit.edu/cgi-bin/primer/primer3.cgi](http://www-genome.wi.mit.edu/cgi-bin/primer/primer3.cgi)), STSPipeline ([www-genome.wi.mit.edu/cgi-bin/www-STSPipeline](http://www-genome.wi.mit.edu/cgi-bin/www-STSPipeline)), or GeneUp (Pesole *et al.*, *BioTechniques* 25:112-123 (1998)), for  
15 example, can be used to identify potential PCR primers.

Nucleic acid molecules or fragments thereof of the present invention are capable of specifically hybridizing to other nucleic acid molecules under certain circumstances. Nucleic  
20 acid molecules of the present invention include those that specifically hybridize to nucleic acid molecules having a nucleic acid sequence selected from the group consisting of SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, complements thereof, and fragments of either. As used herein, two nucleic acid molecules are said to be capable of specifically hybridizing to one another if the two  
25 molecules are capable of forming an anti-parallel, double-stranded nucleic acid structure.

A nucleic acid molecule of the invention can also encode a homolog nucleic acid molecule. As used herein, a homolog nucleic acid molecule or fragment thereof is a counterpart nucleic acid molecule or fragment thereof in a second species (*e.g.*, corn *FATB* intron I nucleic acid molecule is a homolog of *Arabidopsis* *FATB* intron I nucleic acid  
30 molecule). A homolog can also be generated by molecular evolution or DNA shuffling techniques, so that the molecule retains at least one functional or structure characteristic of the original polypeptide (*see*, for example, U.S. Patent 5,811,238).

In another embodiment, the homolog is obtained from a plant selected from the group consisting of alfalfa, *Arabidopsis*, barley, *Brassica campestris*, oilseed rape, broccoli, cabbage, canola, citrus, cotton, garlic, oat, *Allium*, flax, an ornamental plant, jojoba, corn, peanut, pepper, potato, rapeseed, rice, rye, sorghum, strawberry, sugarcane, sugarbeet, tomato, wheat, poplar, pine, fir, eucalyptus, apple, lettuce, lentils, grape, banana, tea, turf grasses, sunflower, *Phaseolus*, crambe, mustard, castor bean, sesame, cottonseed, linseed, safflower, and oil palm. More particularly, a preferred homolog is obtained from a plant selected from the group consisting of canola, corn, *Brassica campestris*, oilseed rape, soybean, crambe, mustard, castor bean, peanut, sesame, cottonseed, linseed, rapeseed, safflower, oil palm, flax, and sunflower. In an even more preferred embodiment, the homolog is obtained from a plant selected from the group consisting of canola, rapeseed, corn, *Brassica campestris*, oilseed rape, soybean, sunflower, safflower, oil palm, and peanut.

In further embodiment, additional *FATB* introns may be obtained by any method by which additional introns may be identified. In a preferred embodiment, additional soybean *FATB* introns may be obtained by screening a soybean genomic library with a probe of either known exon or intron sequences of soybean *FATB*. The soybean *FATB* gene may then be cloned. In another preferred embodiment, additional soybean *FATB* introns may be obtained by a comparison between soybean genomic sequence and soybean cDNA sequence that allows identification of additional introns. In a more preferred embodiment, additional soybean *FATB* introns may be obtained by screening a soybean genomic library with a probe of either known exon or intron sequences of soybean *FATB*. The soybean *FATB* gene may then be cloned and confirmed and any additional introns may be identified by a comparison between soybean genomic sequence and soybean cDNA sequence. Additional introns may, for example without limitation, be amplified by PCR and used in an embodiment of the present invention.

In another preferred embodiment, an intron, such as for example, a soybean intron, may be cloned by alignment to an intron from another organism, such as, for example, *Arabidopsis*. In this embodiment, the location of an intron, for example, in an *Arabidopsis* amino acid sequence is identified. The *Arabidopsis* amino acid sequence, for example, may then be aligned, for example, with the soybean amino acid sequence, providing a prediction for the location, for example, of additional soybean *FATB* introns. Primers may be synthesized, for example, using the soybean *FATB* cDNA. The predicted introns may be

synthesized, for example by PCR, using such primers. Such introns may be used in an embodiment of the present invention.

#### Plant Constructs and Plant Transformants

One or more of the nucleic acid molecules of the invention may be used in plant transformation or transfection. Exogenous genetic material may be transferred into a plant cell and the plant cell regenerated into a whole, fertile or sterile plant or plant part. Exogenous genetic material is any genetic material, whether naturally occurring or otherwise, from any source that is capable of being inserted into any organism.

In one embodiment of the invention, the expression level of a protein or transcript of one *FATB* gene family member is selectively reduced while leaving the level of a protein or transcript of a second *FATB* gene family member partially unaffected. In a preferred embodiment of the invention, the expression level of a protein or transcript in one *FATB* gene family member is selectively reduced while leaving the level of a protein or transcript of a second *FATB* gene family member substantially unaffected. In a highly preferred embodiment of the invention, the expression level of a protein or transcript of one *FATB* gene family member is selectively reduced while leaving the level of a protein or transcript of a second gene family member essentially unaffected.

As used herein, "partially unaffected" refers to a level of an agent such as a protein or mRNA transcript in which the level of the agent that is partially unaffected is within 80%, more preferably within 60%, and even more preferably within 50% of the level at which it is found in a cell or organism that lacks a nucleic acid molecule capable of selectively reducing another agent.

As used herein, "substantially unaffected" refers to a level of an agent such as a protein or mRNA transcript in which the level of the agent that is substantially unaffected is within 49%, more preferably within 35%, and even more preferably within 24% of the level at which it is found in a cell or organism that lacks a nucleic acid molecule capable of selectively reducing another agent.

As used herein, "essentially unaffected" refers to a level of an agent such as a protein or mRNA transcript that is either not altered by a particular event or altered only to an extent that does not affect the physiological function of that agent. In a preferred aspect, the level of an agent that is essentially unaffected is within 20%, more preferably within 10%, and even more preferably within 5% of the level at which it is found in a cell or organism that lacks a nucleic acid molecule capable of selectively reducing another agent.

In a more particularly preferred embodiment, a soybean plant of the present invention includes nucleic acid sequences which when expressed are capable of selectively reducing the expression level of a *FATB* protein and/or *FATB* transcript while overexpressing the level of a different protein and/or transcript. Preferably, the protein is selected from the group consisting of beta-ketoacyl-ACP synthase I, beta-ketoacyl-ACP synthase IV, and delta-9 desaturase, and the different transcript encodes an enzyme selected from the group consisting of beta-ketoacyl-ACP synthase I, beta-ketoacyl-ACP synthase IV, and delta-9 desaturase.

In embodiments where nucleic acid sequences which when expressed in a transformed plant are capable of selectively reducing the expression level of a *FATB* protein and/or *FATB* transcript, preferred nucleic acid sequences are selected from the groups consisting of (1) nucleic acid sequences with at least 50%, 60%, 70%, 80%, 85%, 90%, 95%, 97%, 98%, 99% or 100% sequence identity over the entire length of the nucleic acid molecule with a nucleotide sequence selected from the group consisting of SEQ ID NO: 2, SEQ ID NO:3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, complements thereof, and fragments of either; (2) nucleic acid molecules which contain sequences that are also found in a soybean *FATB* gene intron; and (3) nucleic acid molecules that exhibit sequences with at least 50%, 60%, 70%, 80%, 85%, 90%, 95%, 97%, 98%, 99% or 100% sequence identity over the entire length of the nucleic acid molecule with a nucleic acid molecule of (2).

In a preferred embodiment, a soybean seed of the present invention has an oil composition that is 50% or greater oleic acid and 15% or less saturated fatty acids (including palmitic acid and stearic acid). In a more preferred embodiment, a soybean seed of the present invention has an oil composition that is 10% or less saturated fatty acids. As used herein, all % composition of oils within a plant or plant part such as a seed are determined by weight.

In a particularly preferred embodiment, a soybean seed of the present invention has an oil composition that is 9% or less, 8% or less, 7% or less, 6% or less, 5% or less, 4% or less, 3.6% or less, 3.5% or less, or 3.4% or less saturated fatty acids. In a more preferred embodiment, a soybean seed of the present invention has an oil composition that is a low saturate composition, and in another more preferred embodiment, a soybean seed of the present invention has an oil composition that is a zero saturate composition.

In another preferred embodiment a soybean seed of the present invention has an oil composition that is 50% or greater oleic acid, and between 10 and 15% saturated fatty acids.

In a more preferred embodiment, a soybean seed of the present invention has an oil composition that is between 7 and 10% saturated fatty acids, between 5 and 8% saturated fatty acids, between 3.4 and 7% saturated fatty acids, between 3.5 and 7% saturated fatty acids, between 3.6 and 7% saturated fatty acids, between 2 and 4% saturated fatty acids, or less than 3.4% saturated fatty acids.

In another preferred embodiment of the present invention, a soybean seed of the present invention has an oil composition in which the level of palmitic acid is at least partially reduced, at least substantially reduced, or effectively eliminated. In another embodiment, a soybean seed of the present invention has an oil composition in which the level of stearic acid is at least partially reduced, at least substantially reduced, or effectively eliminated.

In embodiments where nucleic acid sequences which when expressed are capable of selectively reducing the expression level of a *FATB* protein and/or *FATB* transcript such that a soybean seed of the present invention has a low saturate or zero saturate oil composition that also contains 50% or greater oleic acid, the nucleic acid sequences are selected from the groups consisting of: (1) nucleic acid sequences with at least 50%, 60%, 70%, 80%, 85%, 90%, 95%, 97%, 98%, 99% or 100% sequence identity over the entire length of the nucleic acid molecule with a nucleotide sequence selected from the group consisting of SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, complements thereof, and fragments of either; (2) nucleic acid molecules which contain sequences that are also found in a soybean *FATB* intron; and (3) nucleic acid molecules that exhibit sequences with at least 50%, 60%, 70%, 80%, 85%, 90%, 95%, 97%, 98%, 99% or 100% sequence identity over the entire length of the nucleic acid molecule with a nucleic acid molecule of (2).

Genetic material may also be obtained from other species, for example monocotyledons or dicotyledons, including, but not limited to canola, corn, soybean, *Arabidopsis*, *Phaseolus*, peanut, alfalfa, wheat, rice, oat, sorghum, rapeseed, rye, barley, millet, fescue, perennial ryegrass, sugarcane, cranberry, papaya, banana, safflower, oil palm, flax, muskmelon, apple, cucumber, dendrobium, gladiolus, chrysanthemum, liliacea, cotton, eucalyptus, sunflower, *Brassica campestris*, oilseed rape, turfgrass, sugarbeet, coffee and dioscorea (Christou, INO: *Particle Bombardment for Genetic Engineering of Plants*, Biotechnology Intelligence Unit. Academic Press, San Diego, California (1996)), with canola, corn, *Brassica campestris*, oilseed rape, rapeseed, soybean, crambe, mustard, castor

bean, peanut, sesame, cottonseed, linseed, safflower, oil palm, flax, and sunflower preferred, and canola, rapeseed, corn, *Brassica campestris*, oilseed rape, soybean, sunflower, safflower, oil palms, and peanut more preferred. In a more preferred embodiment, canola genetic material is transferred into canola. In another more preferred embodiment, oilseed rape genetic material is transferred into oilseed rape. In another particularly preferred embodiment, soybean genetic material is transferred into soybean.

The levels of products such as transcripts or proteins may be increased or decreased throughout an organism such as a plant or localized in one or more specific organs or tissues of the organism. For example the levels of products may be increased or decreased in one or more of the tissues and organs of a plant including without limitation: roots, tubers, stems, leaves, stalks, fruit, berries, nuts, bark, pods, seeds and flowers. A preferred organ is a seed.

Exogenous genetic material may be transferred into a host cell by the use of a DNA vector or construct designed for such a purpose. Design of such a vector is generally within the skill of the art (See, e.g., *Plant Molecular Biology: A Laboratory Manual*, Clark (ed.), Springer, New York (1997)).

A construct or vector may include a plant promoter to express a nucleic acid molecule of choice. In a preferred embodiment, any nucleic acid molecules described herein can be operably linked to a promoter region which functions in a plant cell to cause the production of an mRNA molecule. For example, any promoter that functions in a plant cell to cause the production of an mRNA molecule, such as those promoters described herein, without limitation, can be used. In a preferred embodiment, the promoter is a plant promoter.

A number of promoters that are active in plant cells have been described in the literature. These include, but are not limited to, the nopaline synthase (NOS) promoter (Ebert *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 84:5745-5749 (1987)), the octopine synthase (OCS) promoter (which is carried on tumor-inducing plasmids of *Agrobacterium tumefaciens*), the caulimovirus promoters such as the cauliflower mosaic virus (CaMV) 19S promoter (Lawton *et al.*, *Plant Mol. Biol.* 9:315-324 (1987)) and the CaMV 35S promoter (Odell *et al.*, *Nature* 313:810-812 (1985)), the figwort mosaic virus 35S-promoter (U.S. Patent No. 5,378,619), the light-inducible promoter from the small subunit of ribulose-1,5-bis-phosphate carboxylase (ssRUBISCO), the Adh promoter (Walker *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 84:6624-6628 (1987)), the sucrose synthase promoter (Yang *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 87:4144-4148 (1990)), the R gene complex promoter (Chandler *et al.*, *The Plant Cell* 1:1175-1183 (1989)) and the chlorophyll a/b binding protein gene promoter. These promoters have

been used to create DNA constructs that have been expressed in plants; *see, e.g.*, PCT publication WO 84/02913. The CaMV 35S promoters are preferred for use in plants. Promoters known or found to cause transcription of DNA in plant cells can be used in the invention.

5           Particularly preferred promoters can also be used to express a nucleic acid molecule of the present invention in seeds or fruits. Indeed, in a preferred embodiment, the promoter used is a seed specific promoter. Examples of such promoters include the 5' regulatory regions from such genes as napin (Kridl *et al.*, *Seed Sci. Res.* 1:209-219 (1991)), phaseolin (Bustos *et al.*, *Plant Cell*, 1(9):839-853 (1989)), soybean trypsin inhibitor (Riggs *et al.*, *Plant*  
10 *Cell* 1(6):609-621 (1989)), ACP (Baerson *et al.*, *Plant Mol. Biol.*, 22(2):255-267 (1993)), stearyl-ACP desaturase (Slocombe *et al.*, *Plant Physiol.* 104(4):167-176 (1994)), soybean  $\alpha'$  subunit of b-conglycinin (soy 7s, (Chen *et al.*, *Proc. Natl. Acad. Sci.*, 83:8560-8564 (1986))), and oleosin (*see, for example*, Hong *et al.*, *Plant Mol. Biol.*, 34(3):549-555 (1997)). Further examples include the promoter for  $\beta$ -conglycinin (Chen *et al.*, *Dev. Genet.* 10: 112-122  
15 (1989)) and the promoter for FAE (PCT Publication WO 01/11061). Preferred promoters for expression in the seed are 7S and napin promoters.

Additional promoters that may be utilized are described, for example, in U.S. Patents 5,378,619; 5,391,725; 5,428,147; 5,447,858; 5,608,144; 5,608,144; 5,614,399; 5,633,441; 5,633,435; and 4,633,436. In addition, a tissue specific enhancer may be used (Fromm *et al.*,  
20 *The Plant Cell* 1:977-984 (1989)).

Constructs or vectors may also include, with the region of interest, a nucleic acid sequence that acts, in whole or in part, to terminate transcription of that region. A number of such sequences have been isolated, including the Tr7 3' sequence and the NOS 3' sequence (Ingelbrecht *et al.*, *The Plant Cell* 1:671-680 (1989); Bevan *et al.*, *Nucleic Acids Res.* 11:369-  
25 385 (1983)). Regulatory transcript termination regions can be provided in plant expression constructs of this invention as well. Transcript termination regions can be provided by the DNA sequence encoding the gene of interest or a convenient transcription termination region derived from a different gene source, for example, the transcript termination region that is naturally associated with the transcript initiation region. The skilled artisan will recognize  
30 that any convenient transcript termination region that is capable of terminating transcription in a plant cell can be employed in the constructs of the present invention.

A vector or construct may also include regulatory elements. Examples of such include the Adh intron 1 (Callis *et al.*, *Genes and Develop.* 1:1183-1200 (1987)), the sucrose

synthase intron (Vasil *et al.*, *Plant Physiol.* 91:1575-1579 (1989)) and the TMV omega element (Gallie *et al.*, *The Plant Cell* 1:301-311 (1989)). These and other regulatory elements may be included when appropriate.

A vector or construct may also include a selectable marker. Selectable markers may also be used to select for plants or plant cells that contain the exogenous genetic material. Examples of such include, but are not limited to: a *neo* gene (Potrykus *et al.*, *Mol. Gen. Genet.* 199:183-188 (1985)), which codes for kanamycin resistance and can be selected for using kanamycin, RptII, G418, hpt; a bar gene which codes for bialaphos resistance; a mutant EPSP synthase gene (Hinchee *et al.*, *Bio/Technology* 6:915-922 (1988); Reynaerts *et al.*,  
10 Selectable and Screenable Markers. In Gelvin and Schilperoort. *Plant Molecular Biology Manual*, Kluwer, Dordrecht (1988); Reynaerts *et al.*, *Selectable and screenable markers*. In Gelvin and Schilperoort. *Plant Molecular Biology Manual*, Kluwer, Dordrecht (1988)), *aadA* (Jones *et al.*, *Mol. Gen. Genet.* (1987)), which encodes glyphosate resistance; a nitrilase gene which confers resistance to bromoxynil (Stalker *et al.*, *J. Biol. Chem.* 263:6310-6314 (1988));  
15 a mutant acetolactate synthase gene (ALS) which confers imidazolinone or sulphonylurea resistance (European Patent Application 154,204 (Sept. 11, 1985)), ALS (D'Halluin *et al.*, *Bio/Technology* 10: 309-314 (1992)), and a methotrexate resistant DHFR gene (Thillet *et al.*, *J. Biol. Chem.* 263:12500-12508 (1988)).

A vector or construct may also include a screenable marker. Screenable markers may be used to monitor expression. Exemplary screenable markers include: a  $\beta$ -glucuronidase or  
20 *uidA* gene (GUS) which encodes an enzyme for which various chromogenic substrates are known (Jefferson, *Plant Mol. Biol. Rep.* 5:387-405 (1987); Jefferson *et al.*, *EMBO J.* 6:3901-3907 (1987)); an R-locus gene, which encodes a product that regulates the production of anthocyanin pigments (red color) in plant tissues (Dellaporta *et al.*, *Stadler Symposium*  
25 11:263-282 (1988)); a  $\beta$ -lactamase gene (Sutcliffe *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 75:3737-3741 (1978)), a gene which encodes an enzyme for which various chromogenic substrates are known (*e.g.*, PADAC, a chromogenic cephalosporin); a luciferase gene (Ow *et al.*, *Science* 234:856-859 (1986)); a *xylE* gene (Zukowsky *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 80:1101-1105 (1983)) which encodes a catechol dioxygenase that can convert  
30 chromogenic catechols; an  $\alpha$ -amylase gene (Ikata *et al.*, *Bio/Technol.* 8:241-242 (1990)); a tyrosinase gene (Katz *et al.*, *J. Gen. Microbiol.* 129:2703-2714 (1983)) which encodes an enzyme capable of oxidizing tyrosine to DOPA and dopaquinone which in turn condenses to melanin; an  $\alpha$ -galactosidase, which will turn a chromogenic  $\alpha$ -galactose substrate.

Included within the terms "selectable or screenable marker genes" are also genes that encode a secretable marker whose secretion can be detected as a means of identifying or selecting for transformed cells. Examples include markers that encode a secretable antigen that can be identified by antibody interaction, or even secretable enzymes that can be detected catalytically. Secretable proteins fall into a number of classes, including small, diffusible proteins that are detectable, (*e.g.*, by ELISA), small active enzymes that are detectable in extracellular solution (*e.g.*,  $\alpha$ -amylase,  $\beta$ -lactamase, phosphinothricin transferase), or proteins that are inserted or trapped in the cell wall (such as proteins that include a leader sequence such as that found in the expression unit of extension or tobacco PR-S). Other possible selectable and/or screenable marker genes will be apparent to those of skill in the art.

It is understood that two or more nucleic molecules of the present invention may be introduced into a plant using a single construct and that construct can contain more than one promoter. In embodiments where the construct is designed to express two nucleic acid molecules, it is preferred that the two promoters are (i) two constitutive promoters, (ii) two seed-specific promoters, or (iii) one constitutive promoter and one seed-specific promoter. Preferred seed-specific and constitutive promoters are a napin and a 7S promoter, respectively. Illustrative combinations are set forth in Example 5. It is understood that two or more of the nucleic molecules may be physically linked and expressed utilizing a single promoter, preferably a seed-specific or constitutive promoter.

It is further understood that two or more nucleic acids of the present invention may be introduced into a plant using two or more different constructs. Alternatively, two or more nucleic acids of the present invention may be introduced into two different plants and the plants may be crossed to generate a single plant expressing two or more nucleic acids. In an RNAi embodiment, it is understood that the sense and antisense strands may be introduced into the same plant on one construct or two constructs. Alternatively, the sense and antisense strands may be introduced into two different plants and the plants may be crossed to generate a single plant expressing both sense and antisense strands.

Any of the nucleic acid molecules and constructs of the invention may be introduced into a plant or plant cell in a permanent or transient manner. Preferred nucleic acid molecules and constructs of the present invention are described above in the Detailed Description, and in the Examples. Another embodiment of the invention is directed to a method of producing transgenic plants which generally comprises the steps of selecting a suitable plant or plant

cell, transforming the plant or plant cell with a recombinant vector, and obtaining a transformed host cell.

In a preferred embodiment the plant or cell is, or is derived from, a plant involved in the production of vegetable oils for edible and industrial uses. Especially preferred are temperate oilseed crops. Plants of interest include, but are not limited to, rapeseed (canola and High Erucic Acid varieties), maize, soybean, crambe, mustard, castor bean, peanut, sesame, cotton, linseed, safflower, oil palm, flax, sunflower, and coconut. The invention is applicable to monocotyledonous or dicotyledonous species alike, and will be readily applicable to new and/or improved transformation and regulatory techniques.

Methods and technology for introduction of DNA into plant cells are well known to those of skill in the art, and virtually any method by which nucleic acid molecules may be introduced into a cell is suitable for use in the present invention. Non-limiting examples of suitable methods include: chemical methods; physical methods such as microinjection, electroporation, the gene gun, microprojectile bombardment, and vacuum infiltration; viral vectors; and receptor-mediated mechanisms. Other methods of cell transformation can also be used and include but are not limited to introduction of DNA into plants by direct DNA transfer into pollen, by direct injection of DNA into reproductive organs of a plant, or by direct injection of DNA into the cells of immature embryos followed by the rehydration of desiccated embryos.

*Agrobacterium*-mediated transfer is a widely applicable system for introducing genes into plant cells. See, e.g., Fraley *et al.*, *Bio/Technology* 3:629-635 (1985); Rogers *et al.*, *Methods Enzymol.* 153:253-277 (1987). The region of DNA to be transferred is defined by the border sequences and intervening DNA is usually inserted into the plant genome. Spielmann *et al.*, *Mol. Gen. Genet.* 205:34 (1986). Modern *Agrobacterium* transformation vectors are capable of replication in *E. coli* as well as *Agrobacterium*, allowing for convenient manipulations. Klee *et al.*, In: *Plant DNA Infectious Agents*, Hohn and Schell (eds.), Springer-Verlag, New York, pp. 179-203 (1985).

The regeneration, development and cultivation of plants from single plant protoplast transformants or from various transformed explants is well known in the art. See generally, Maliga *et al.*, *Methods in Plant Molecular Biology*, Cold Spring Harbor Press (1995); Weissbach and Weissbach, In: *Methods for Plant Molecular Biology*, Academic Press, San Diego, CA (1988). Plants of the present invention can be part of or generated from a

breeding program, and may also be reproduced using apomixis. Methods for the production of apomictic plants are known in the art. See, e.g., U.S. Patent 5,811,636.

Cosuppression is the reduction in expression levels, usually at the level of RNA, of a particular endogenous gene or gene family by the expression of a homologous sense  
5 construct that is capable of transcribing mRNA of the same strandedness as the transcript of the endogenous gene (Napoli *et al.*, *Plant Cell* 2:279-289 (1990); van der Krol *et al.*, *Plant Cell* 2:291-299 (1990)). Cosuppression may result from stable transformation with a single copy nucleic acid molecule that is homologous to a nucleic acid sequence found with the cell (Prolls and Meyer, *Plant J.* 2:465-475 (1992)) or with multiple copies of a nucleic acid  
10 molecule that is homologous to a nucleic acid sequence found with the cell (Mittlesten *et al.*, *Mol. Gen. Genet.* 244:325-330 (1994)). Genes, even though different, linked to homologous promoters may result in the cosuppression of the linked genes (Vaucheret, *C.R. Acad. Sci. III* 316:1471-1483 (1993); Flavell, *Proc. Natl. Acad. Sci. (U.S.A.)* 91:3490-3496 (1994)); van Blokland *et al.*, *Plant J.* 6:861-877 (1994); Jorgensen, *Trends Biotechnol.* 8:340-344 (1990);  
15 Meins and Kunz, In: *Gene Inactivation and Homologous Recombination in Plants*, Paszkowski (ed.), pp. 335-348, Kluwer Academic, Netherlands (1994))(Kinney, Induced Mutations and Molecular Techniques for Crop Improvement, Proceedings of a Symposium 19-23 June 1995 (jointly organized by IAEA and FA)), pages 101-113 (IAEA-SM 340-49).

It is understood that one or more of the nucleic acids of the invention may be  
20 introduced into a plant cell and transcribed using an appropriate promoter with such transcription resulting in the cosuppression of an endogenous protein.

Antisense approaches are a way of preventing or reducing gene function by targeting the genetic material (Mol *et al.*, *FEBS Lett.* 268:427-430 (1990)). The objective of the antisense approach is to use a sequence complementary to the target gene to block its  
25 expression and create a mutant cell line or organism in which the level of a single chosen protein is selectively reduced or abolished. Antisense techniques have several advantages over other 'reverse genetic' approaches. The site of inactivation and its developmental effect can be manipulated by the choice of promoter for antisense genes or by the timing of external application or microinjection. Antisense can manipulate its specificity by selecting either  
30 unique regions of the target gene or regions where it shares homology to other related genes (Hiatt *et al.*, In: *Genetic Engineering*, Setlow (ed.), Vol. 11, New York: Plenum 49-63 (1989)).

Antisense RNA techniques involve introduction of RNA that is complementary to the target mRNA into cells, which results in specific RNA:RNA duplexes being formed by base pairing between the antisense substrate and the target mRNA (Green *et al.*, *Annu. Rev. Biochem.* 55:569-597 (1986)). Under one embodiment, the process involves the introduction and expression of an antisense gene sequence. Such a sequence is one in which part or all of the normal gene sequences are placed under a promoter in inverted orientation so that the 'wrong' or complementary strand is transcribed into a noncoding antisense RNA that hybridizes with the target mRNA and interferes with its expression (Takayama and Inouye, *Crit. Rev. Biochem. Mol. Biol.* 25:155-184 (1990)). An antisense vector is constructed by standard procedures and introduced into cells by methods including but not limited to transformation, transfection, electroporation, microinjection, and infection. The type of transformation and choice of vector will determine whether expression is transient or stable. The promoter used for the antisense gene may influence the level, timing, tissue, specificity, or inducibility of the antisense inhibition.

It has been reported that the introduction of double-stranded RNA into cells may also be used to disrupt the function of an endogenous gene. (Fire *et al.*, *Nature* 391: 806-811 (1998)). Such disruption has been demonstrated, for example, in *Caenorhabditis elegans* and is often referred to as RNA interference, or RNAi. (Fire *et al.*, *Nature* 391: 806-811 (1998)). The disruption of gene expression in *C. elegans* by double-stranded RNA has been reported to induce suppression by a post-transcriptional mechanism. (Montgomery *et al.*, *Proc. Natl. Acad. Sci.* 95:15502-15507 (1998)). Evidence of gene silencing by double-stranded RNA has also been reported for plants. (Waterhouse *et al.*, *Proc. Natl. Acad. Sci.* 95: 13959-13964 (1998)). See also Plasterk, *Science* 296: 1263-1265 (2002); Zamore, *Science* 296: 1265-1269 (2002).

An intron-spliced hairpin structure reportedly may also be used to effect post-transcriptional gene suppression. (Smith *et al.*, *Nature* 407: 319-320 (2000)). Reports indicate that post-transcriptional gene silencing can be induced with almost 100% efficiency by the use of intron-spliced RNA with a hairpin structure. (Smith *et al.*, *Nature* 407: 319-320 (2000)). Representative methods for effecting RNA silencing are set forth in U.S. Application, Attorney Docket No. 16518.069, entitled "Intron Double Stranded RNA Constructs And Uses Thereof," naming JoAnne Fillatti as inventor, filed concurrently herewith.

It is understood that one or more of the nucleic acids of the invention may be modified in order to effect RNAi or another mode of post-transcriptional gene suppression.

The present invention also provides for parts of the plants, particularly reproductive or storage parts. Plant parts, without limitation, include seed, endosperm, ovule, pollen, roots, tubers, stems, leaves, stalks, fruit, berries, nuts, bark, pods, seeds and flowers. In a particularly preferred embodiment of the present invention, the plant part is a seed.

The present invention also provides a container of over 10,000, more preferably 20,000, and even more preferably 40,000 seeds where over 10%, more preferably 25%, more preferably 50% and even more preferably 75% or 90% of the seeds are seeds derived from a plant of the present invention.

The present invention also provides a container of over 10 kg, more preferably 25 kg, and even more preferably 50 kg seeds where over 10%, more preferably 25%, more preferably 50% and even more preferably 75% or 90% of the seeds are seeds derived from a plant of the present invention.

Any of the plants or parts thereof of the present invention may be processed to produce a feed, meal, protein, or oil preparation. A particularly preferred plant part for this purpose is a seed. In a preferred embodiment the feed, meal, protein or oil preparation is designed for livestock animals or humans, or both. Methods to produce feed, meal, protein and oil preparations are known in the art. See, for example, U.S. Patents 4,957,748, 5,100,679, 5,219,596, 5,936,069, 6,005,076, 6,146,669, and 6,156,227. In a preferred embodiment, the protein preparation is a high protein preparation. Such a high protein preparation preferably has a protein content of greater than 5% w/v, more preferably 10% w/v, and even more preferably 15% w/v. In a preferred oil preparation, the oil preparation is a high oil preparation with an oil content derived from a plant or part thereof of the present invention of greater than 5% w/v, more preferably 10% w/v, and even more preferably 15% w/v. In a preferred embodiment the oil preparation is a liquid and of a volume greater than 1, 5, 10 or 50 liters. The present invention provides for oil produced from plants of the present invention or generated by a method of the present invention. Such an oil may exhibit enhanced oxidative stability. Also, such oil may be a minor or major component of any resultant product. Moreover, such oil may be blended with other oils. In a preferred embodiment, the oil produced from plants of the present invention or generated by a method of the present invention constitutes greater than 0.5%, 1%, 5%, 10%, 25%, 50%, 75% or 90% by volume or weight of the oil component of any product. In another embodiment, the oil

preparation may be blended and can constitute greater than 10%, 25%, 35%, 50% or 75% of the blend by volume. Oil produced from a plant of the present invention can be admixed with one or more organic solvents or petroleum distillates.

In one embodiment, an oil of the present invention has an oil composition that is 50% or greater oleic acid and 15% or less saturated fatty acids. In another embodiment, an oil of the present invention has an oil composition that is 10% or less saturated fatty acids. In another embodiment, an oil of the present invention has an oil composition that is 9% or less saturated fatty acids, 8% or less saturated fatty acids, 7% or less saturated fatty acids, 6% or less saturated fatty acids, 5% or less saturated fatty acids, 4% or less saturated fatty acids, 3.6% or less saturated fatty acids, 3.5% or less saturated fatty acids, or 3.4% or less saturated fatty acids. In a more preferred embodiment, an oil of the present invention has a low saturate oil composition, and in another preferred embodiment, an oil of the present invention has a zero saturate oil composition.

In another preferred embodiment, an oil of the present invention has an oil composition that is 50% or greater oleic acid, and between 10 and 15% saturated fatty acids. In a more preferred embodiment, an oil of the present invention has an oil composition that is between 7 and 10% saturated fatty acids, between 5 and 8% saturated fatty acids, between 3.4 and 7% saturated fatty acids, between 3.5 and 7% saturated fatty acids, between 3.6 and 7% saturated fatty acids, between 2 and 4% saturated fatty acids, or less than 3.4% saturated fatty acids.

In another preferred embodiment, an oil of the present invention has an oil composition in which the level of palmitic acid is at least partially reduced, at least substantially reduced, or effectively eliminated. In another embodiment, an oil of the present invention has an oil composition in which the level of stearic acid is at least partially reduced, at least substantially reduced, or effectively eliminated.

In embodiments where nucleic acid sequences which when expressed are capable of selectively reducing the expression level of a protein and/or transcript encoded by a *FATB* gene such that an oil of the present invention has an oil composition that is 50% or greater oleic acid, and 10% or less saturated fatty acids, preferably 5% or less saturated fatty acids, preferably 3.6% or less saturated fatty acids, preferably 3.5% or less saturated fatty acids, and more preferably 3.4% or less saturated fatty acids, the nucleic acid sequences are selected from the groups consisting of: (1) nucleic acid sequences with at least 50%, 60%, 70%, 80%, 85%, 90%, 95%, 97%, 98%, 99% or 100% sequence identity over the entire length of the

nucleic acid molecule with a nucleotide sequence selected from the group consisting of SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, complements thereof, and fragments of either; (2) nucleic acid molecules which contain sequences that are also found in a soybean *FATB* gene intron; and (3) nucleic acid molecules that exhibit sequences with at least 50%, 60%, 70%, 80%, 85%, 90%, 95%, 97%, 98%, 99% or 100% sequence identity over the entire length of the nucleic acid molecule with a nucleic acid molecule of (2).

#### Computer Readable Medium

The nucleotide sequence provided in SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, complements thereof, and fragments of either, or a nucleotide sequence at least 50%, 60%, or 70% identical, preferably 80%, 85% identical, or especially preferably 90%, or 95% identical, or particularly highly preferably 97%, 98%, or 99% identical to the sequence provided in SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, complements thereof, and fragments of either, can be "provided" in a variety of media to facilitate use. Such a medium can also provide a subset thereof in a form that allows a skilled artisan to examine the sequences.

In one application of this embodiment, a nucleotide sequence of the present invention can be recorded on computer readable media. As used herein, "computer readable media" refers to any medium that can be read and accessed directly by a computer. Such media include, but are not limited to: magnetic storage media, such as floppy disk, hard disk, storage medium, and magnetic tape; optical storage media such as CD-ROM; electrical storage media such as RAM and ROM; and hybrids of these categories such as magnetic/optical storage media. A skilled artisan can readily appreciate how any of the presently known computer readable media can be used to create a manufacture comprising a computer readable medium having recorded thereon a nucleotide sequence of the present invention.

As used herein, "recorded" refers to a process for storing information on computer readable medium. A skilled artisan can readily adopt any of the presently known methods for recording information on computer readable media to generate media comprising the nucleotide sequence information of the present invention. A variety of data storage structures are available to a skilled artisan for creating a computer readable medium having recorded thereon a nucleotide sequence of the present invention. The choice of the data storage

structure will generally be based on the means chosen to access the stored information. In addition, a variety of data processor programs and formats can be used to store the nucleotide sequence information of the present invention on computer readable media. The sequence information can be represented in a word processing text file, formatted in commercially-  
5 available software such as Word Perfect and Microsoft Word, or represented in the form of an ASCII file, stored in a database application, such as DB2, Sybase, Oracle, or the like. A skilled artisan can readily adapt any number of data processor structuring formats (*e.g.*, text file or database) in order to obtain computer readable media having recorded thereon the nucleotide sequence information of the present invention.

10 By providing one or more nucleotide sequences of the present invention, a skilled artisan can routinely access the sequence information for a variety of purposes. Computer software is publicly available which allows a skilled artisan to access sequence information provided in a computer readable medium. Software which implements the BLAST (Altschul *et al.*, *J. Mol. Biol.* 215: 403-410 (1990)) and BLAZE (Brutlag *et al.*, *Comp. Chem.* 17:203-  
15 207 (1993)) search algorithms on a Sybase system can be used to identify non-coding regions and other nucleic acid molecules of the present invention within the genome that contain homology to non-coding regions from other organisms. Such non-coding regions may be utilized to affect the expression of commercially important proteins such as enzymes used in amino acid biosynthesis, metabolism, transcription, translation, RNA processing, nucleic acid  
20 and protein degradation, protein modification, and DNA replication, restriction, modification, recombination, and repair.

The present invention further provides systems, particularly computer-based systems, which contain the sequence information described herein. Such systems are designed to identify commercially important fragments of the nucleic acid molecules of the present  
25 invention. As used herein, "a computer-based system" refers to the hardware means, software means, and data storage means used to analyze the nucleotide sequence information of the present invention. The minimum hardware means of the computer-based systems of the present invention comprises a central processing unit (CPU), input means, output means, and data storage means. A skilled artisan can readily appreciate that any one of the currently  
30 available computer-based systems is suitable for use in the present invention.

As indicated above, the computer-based systems of the present invention comprise a data storage means having stored therein a nucleotide sequence of the present invention and the necessary hardware means and software means for supporting and implementing a search

means. As used herein, "data storage means" refers to memory that can store nucleotide sequence information of the present invention, or a memory access means which can access manufactures having recorded thereon the nucleotide sequence information of the present invention. As used herein, "search means" refers to one or more programs which are  
5 implemented on the computer-based system to compare a target sequence or target structural motif with the sequence information stored within the data storage means. Search means are used to identify fragments or regions of the sequence of the present invention that match a particular target sequence or target motif. A variety of known algorithms are disclosed publicly and a variety of commercially available software for conducting search means are  
10 available and can be used in the computer-based systems of the present invention. Examples of such software include, but are not limited to, MacPattern (EMBL), BLASTIN, and BLASTIX (NCBIA). One of the available algorithms or implementing software packages for conducting homology searches can be adapted for use in the present computer-based systems.

The most preferred sequence length of a target sequence is from about 10 to 100  
15 amino acids or from about 30 to 300 nucleotide residues. However, it is well recognized that during searches for commercially important fragments of the nucleic acid molecules of the present invention, such as sequence fragments involved in gene expression and protein processing, the target sequence may be of shorter length.

As used herein, "a target structural motif," or "target motif" refers to any rationally  
20 selected sequence or combination of sequences in which the sequences are chosen based on a three-dimensional configuration which is formed upon the folding of the target motif. There are a variety of target motifs known in the art. Protein target motifs include, but are not limited to, enzymatic active sites and signal sequences. Nucleic acid target motifs include, but are not limited to, promoter sequences, *cis* elements, hairpin structures, and inducible  
25 expression elements (protein binding sequences).

Thus, the present invention further provides an input means for receiving a target sequence, a data storage means for storing the target sequences of the present invention sequence identified using a search means as described above, and an output means for outputting the identified homologous sequences. A variety of structural formats for the input  
30 and output means can be used to input and output information in the computer-based systems of the present invention. A preferred format for an output means ranks fragments of the sequence of the present invention by varying degrees of homology to the target sequence or target motif. Such presentation provides a skilled artisan with a ranking of sequences which

contain various amounts of the target sequence or target motif and identifies the degree of homology contained in the identified fragment.

A variety of comparing means can be used to compare a target sequence or target motif with the data storage means to identify sequence fragments sequence of the present invention. For example, implementing software that implement the BLAST and BLAZE algorithms (Altschul *et al.*, *J. Mol. Biol.* 215:403-410 (1990)) can be used to identify non-coding regions within the nucleic acid molecules of the present invention. A skilled artisan can readily recognize that any one of the publicly available homology search programs can be used as the search means for the computer-based systems of the present invention.

The following examples are illustrative and not intended to be limiting in any way.

## EXAMPLES

### Example 1 Cloning of *FATB* Thioesterase Genomic Sequences

Leaf tissue is obtained from Asgrow soy variety A3244, ground up in liquid nitrogen and stored at  $-80^{\circ}\text{C}$  until use. Six ml of SDS Extraction buffer (650 ml sterile  $\text{ddH}_2\text{O}$ , 100 ml 1M Tris-Cl pH 8, 100 ml 0.25M EDTA, 50 ml 20% SDS, 100 ml 5M NaCl, 4  $\mu\text{l}$  beta-mercaptoethanol) is added to 2 ml of frozen/ground leaf tissue, and the mixture is incubated at  $65^{\circ}\text{C}$  for 45 minutes. The sample is shaken every 15 minutes. 2 ml of ice-cold 5M potassium acetate is added to the sample, the sample is shaken, and then is incubated on ice for 20 minutes. 3 ml of  $\text{CHCl}_3$  is added to the sample and the sample is shaken for 10 minutes.

The sample is centrifuged at 10,000 rpm for 20 minutes and the supernatant is collected. 2 ml of isopropanol is added to the supernatant and mixed. The sample is then centrifuged at 10,000 rpm for 20 minutes and the supernatant is drained. The pellet is resuspended in 200  $\mu\text{l}$  RNase and incubated at  $65^{\circ}\text{C}$  for 20 minutes. 300  $\mu\text{l}$  ammonium acetate/isopropanol (1:7) is added and mixed. The sample is then centrifuged at 10,000 rpm for 15 minutes and the supernatant was discarded. The pellet is rinsed with 500  $\mu\text{l}$  80% ethanol and allowed to air dry. The pellet of genomic DNA is then resuspended in 200  $\mu\text{l}$  T10E1 (10mM Tris:1mM EDTA).

In a first method, a soy *FATB* cDNA sequence is used to design six oligonucleotides that span the gene: F1 (SEQ ID NO: 11), F2 (SEQ ID NO: 12), F3 (SEQ ID NO: 13), R1 (SEQ ID NO: 14), R2 (SEQ ID NO: 15), and R3 (SEQ ID NO: 16). The oligonucleotides are

used in pairs for PCR amplification from the isolated soy genomic DNA: pair 1 (F1 + R1), pair 2 (F1 + R2), pair 3 (F1 + R3), pair 4 (F2 + R1), pair 5 (F2 + R2), pair 6 (F2 + R3), pair 7 (F3 + R1), and pair 8 (F3 + R2). The PCR amplifications is carried out as follows: 1 cycle, 95°C for 10 minutes; 40 cycles, 95°C for 1 minutes, 58°C for 30 sec, 72°C for 55 sec; 1  
5 cycle, 72°C for 7 minutes. Three positive fragments are obtained, specifically from primer pairs 3, 6, and 7. Each fragment is cloned into vector pCR2.1 (Invitrogen). Cloning is successful only for genomic fragment #3, which is confirmed and sequenced (SEQ ID NO: 10).

Three introns are identified in the soybean *FATB* gene by comparison of the genomic  
10 sequence to the cDNA sequence: intron I (SEQ ID NO: 2) spans base 106 to base 214 of the genomic sequence (SEQ ID NO: 10) and is 109 bp in length; intron II (SEQ ID NO: 3) spans base 289 to base 1125 of the genomic sequence (SEQ ID NO: 10) and is 837 bp in length; and intron III (SEQ ID NO: 4) spans base 1635 to base 1803 of the genomic sequence (SEQ ID NO: 10) and is 169 bp in length.

15 In a second method, the *Arabidopsis thaliana* *FATB* cDNA and *A. thaliana* *FATB* genomic sequence are aligned with the soy *FATB* cDNA and the potential locations of soy *FATB* introns are determined. Oligonucleotides are synthesized for sequences flanking the putative soy introns, and genomic DNA is amplified using appropriate primer pairs. Four additional introns are identified in the soybean *FATB* gene by comparison of the amplified  
20 genomic sequences to the cDNA sequence. These four soy intron sequences are combined with the soy cDNA sequence and the three previously isolated soy intron sequences to generate a genomic sequence of the *FATB* gene (SEQ ID NO: 1). The four new introns isolated are as follows: primers F1 and R1 yield intron IV (SEQ ID NO: 5 ), which spans base 1939 to base 2463 of the genomic sequence (SEQ ID NO: 1) and is 525 bp in length;  
25 primers F2 and R2 yield intron V (SEQ ID NO: 6 ), which spans base 2578 to base 2966 of the genomic sequence (SEQ ID NO: 1) and is 389 bp in length; primers F3 and R3 yield intron VI (SEQ ID NO: 7 ) spans base 3140 to base 3245 of the genomic sequence (SEQ ID NO: 1) and is 106 bp in length and intron VII (SEQ ID NO: 8) which spans base 3314 to base 3395 of the genomic sequence (SEQ ID NO: 1) and is 82 bp in length.

## 30 Example 2 Plant expression constructs

A soybean *FATB* intron II sequence (SEQ ID NO: 3) is amplified via PCR using the partial *FATB* cloned genomic DNA sequence (SEQ ID NO: 10) as a template and primers

18133 (SEQ ID NO: 17) and 18134 (SEQ ID NO: 18). PCR amplification is carried out as follows: 1 cycle, 95°C for 10 minutes; 25 cycles, 95°C for 30 sec, 62°C for 30 sec, 72°C for 30 sec; 1 cycle, 72°C for 7 minutes.

5 PCR amplification resulted in a product (SEQ ID NO: 19) that is 854 bp long. The PCR product is cloned directly into the expression cassette pCGN3892 (Figure 1) in sense orientation, by way of *Xho*I sites engineered onto the 5' ends of the PCR primers, to form pMON70674 (Figure 2). Vector pCGN3892 contains the soybean 7S promoter and a pea RBCS 3'. pMON70674 is then cut with *Nor*I and ligated into pMON41164, a vector that contains the CP4 gene regulated by the FMV promoter (Figure 3). The resulting gene  
10 expression construct, pMON70678 (Figure 4), is used for transformation of soybean using *Agrobacterium* methods as described herein.

Two other expression constructs containing the soybean *FATB* intron II sequence (SEQ ID NO: 3) are created. pMON70674 is cut with *Nor*I and ligated into pMON70675 (Figure 5) which contains the CP4 gene regulated by the FMV promoter and the KAS IV  
15 gene regulated by the napin promoter. The resulting expression construct, pMON70680 (Figure 6), is used for transformation of soybean using *Agrobacterium* methods as described herein. The expression vector pMON70680 is then cut with *Sna*BI and ligated with a gene fusion of the jojoba delta-9 desaturase gene in the sense orientation regulated by the 7S promoter (pMON70656; Figure 7). The resulting expression construct, pMON70681 (Figure  
20 8), is used for transformation of soybean using *Agrobacterium* methods as described herein.

Other soybean *FATB* intron sequences, such as SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, or SEQ ID NO: 8, are cloned in a similar manner. Appropriate primers are designed based on the intron sequence desired. These primer pairs are used to amplify an intron from the *FATB* genomic sequence. The amplified intron is  
25 ligated into the desired expression vector and the construct is transformed into soybean.

### Example 3 Plant Transformation and Analysis

Linear DNA fragments containing the expression constructs of the soybean *FATB* introns are stably introduced into soybean (Asgrow variety A3244) by the method of  
30 Martinell *et al.*, U. S. Patent 6,384,301. Transformed soybean plants are identified by selection on media containing glyphosate.

Fatty acid compositions are analyzed from seed of soybean lines transformed with the intron expression constructs using gas chromatography. R<sub>1</sub> single seed oil compositions of

plants harboring pMON70678 demonstrate that the saturated and unsaturated fatty acid compositions are altered in the oil of seeds from transgenic soybean lines as compared to those of the seed from non-transformed soybean (Table 1). In particular, 16:0 is reduced in transgenic seeds. Selections can be made from such lines depending on the desired relative fatty acid composition. In addition, since each of the introns is able to modify the levels of each fatty acid to varying extents, it is contemplated that combinations of introns can be used depending on the desired compositions.

Table 1

R1 single seed

data

Fatty Acids

Construct	Event	16:0	18:0	18:1	18:2	18:3
PMON70678	GM_A31349	7.7	5.0	17.4	62.2	7.7
PMON70678	GM_A31349	7.8	4.6	18.2	61.9	7.3
PMON70678	GM_A31349	7.9	5.5	17.3	60.4	8.3
PMON70678	GM_A31349	7.9	5.0	17.3	60.6	8.6
PMON70678	GM_A31349	8.2	5.5	15.6	61.8	8.3
PMON70678	GM_A31342	8.4	6.8	12.4	63.1	9.0
PMON70678	GM_A31342	8.7	5.3	15.9	62.7	7.3
PMON70678	GM_A31341	8.7	4.0	19.5	59.4	7.8
PMON70678	GM_A31345	8.8	5.1	15.2	62.4	8.4
PMON70678	GM_A31342	8.8	5.8	13.4	63.0	9.0
PMON70678	GM_A31345	8.9	5.2	15.3	62.0	8.7
PMON70678	GM_A31345	8.9	5.6	15.0	61.9	8.4
PMON70678	GM_A31341	8.9	3.3	38.8	43.2	5.3
PMON70678	GM_A31345	9.0	5.1	16.6	60.7	8.5
PMON70678	GM_A31342	9.0	5.5	16.2	61.9	7.2
PMON70678	GM_A31341	9.0	4.1	31.1	49.9	5.5
PMON70678	GM_A31349	9.1	6.0	12.7	61.9	9.7
PMON70678	GM_A31342	9.1	5.2	15.4	62.5	7.8
PMON70678	GM_A31417	9.2	5.6	15.1	60.8	9.0
PMON70678	GM_A31349	9.2	5.5	14.0	62.2	9.2
PMON70678	GM_A31350	9.2	4.6	18.5	58.8	8.5
PMON70678	GM_A31342	9.4	5.1	15.5	62.2	7.5
PMON70678	GM_A31350	9.5	5.3	14.7	61.5	8.6
PMON70678	GM_A31417	9.5	5.3	15.3	60.9	8.6
PMON70678	GM_A31345	9.5	5.7	14.6	61.2	8.8
PMON70678	GM_A31350	9.6	5.5	13.7	61.7	9.1
PMON70678	GM_A31417	9.6	5.2	16.0	60.0	8.8

PMON70678	GM_A31341	9.6	3.5	24.6	54.9	6.9
PMON70678	GM_A31341	9.7	3.7	20.7	58.5	6.7
PMON70678	GM_A31341	9.8	3.8	19.6	58.5	7.7
PMON70678	GM_A31345	9.9	5.1	14.8	61.4	8.6
control	A3244	12.4	4.3	18.3	56.4	8.0
control	A3244	12.4	6.7	14.0	57.1	8.8
control	A3244	12.6	4.9	15.4	57.4	9.1
control	A3244	12.9	5.0	17.6	55.9	8.2
control	A3244	12.9	4.9	14.4	57.5	9.8
control	A3244	13.0	4.7	14.6	55.6	9.7
control	A3244	13.0	4.7	14.9	57.0	9.4
control	A3244	13.0	5.0	13.8	57.4	10.2
control	A3244	13.2	4.5	16.9	54.6	7.8
control	A3244	13.3	5.1	14.1	57.8	9.4

#### Example 4

RNA is isolated from homozygous R<sub>2</sub> seed from two *FATB* intron suppressed lines, and from negative controls (wild type seed and seed from null segregants from each intron suppressed event). Northern gels containing these RNA samples are probed with the *FATB* cDNA. *FATB* transcript levels are significantly reduced in the intron suppressed lines relative to the negative controls.

#### Example 5 *FATB* Intron Constructs

Plant expression constructs are made containing one or more *FATB* introns in sense or antisense orientation. To achieve a desired fatty acid effect, two or more *FatB* introns are combined into one transcriptional unit. In an alternate approach, each *FATB* intron is expressed under the control of its own promoter (monocistronic). Other constructs are made where a *FATB* intron is capable of producing dsRNA, either using only one transcriptional unit (inverted repeat) or two expression cassettes, with one containing a sense intron and the other containing an antisense intron.

These constructs are stably introduced into soybean (for example, Asgrow variety A3244) by the methods described earlier. Transformed soybean plants are identified by

selection on media containing glyphosate. Gas chromatography is used to determine the fatty acid composition of seed from soybean lines transformed with the constructs. In addition, any of the constructs can contain other sequences of interest, including without limitation, sequences to overexpress KAS I, KAS IV, and/or delta-9 desaturase, as well as different  
5 combinations of promoters.

What is claimed is:

1. A recombinant nucleic acid molecule comprising as operably linked components:  
(A) a promoter that functions in a plant cell to cause production of an mRNA molecule; and  
(B) a nucleic acid sequence that has at least 85% identity to a nucleic acid sequence selected from the group consisting of SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, complements thereof, and fragments of either.
2. The recombinant nucleic acid molecule of claim 1, wherein the promoter is a seed-specific promoter.
3. The recombinant nucleic acid molecule of claim 2, wherein the promoter is a 7S promoter.
4. The recombinant nucleic acid molecule of claim 1, wherein the nucleic acid sequence is in a sense orientation relative to the promoter.
5. The recombinant nucleic acid molecule of claim 1, wherein the nucleic acid sequence is in an antisense orientation relative to the promoter.
6. The recombinant nucleic acid molecule of claim 1, wherein the nucleic acid sequence is capable of expressing a dsRNA.
7. The recombinant nucleic acid molecule of claim 1, wherein said nucleic acid molecule further comprises one or more additional nucleic acid sequences, wherein said additional nucleic acid sequences encode an enzyme selected from the group consisting of beta-ketoacyl-ACP synthase I, beta-ketoacyl-ACP synthase IV, and delta-9 desaturase.
8. The recombinant nucleic acid molecule of claim 7, wherein the additional nucleic acid sequence encodes beta-ketoacyl-ACP synthase IV.
9. The recombinant nucleic acid molecule of claim 7, wherein the additional nucleic acid sequences encode beta-ketoacyl-ACP synthase IV and delta-9 desaturase.

10. An intron obtained from a genomic polynucleotide sequence wherein said genomic polynucleotide sequence is selected from the group consisting of:

- a) a genomic polynucleotide sequence having at least 70% identity to coding regions of SEQ ID NO: 1 over the entire length of SEQ ID NO: 1;
- b) a genomic polynucleotide sequence having at least 80% identity to coding regions of SEQ ID NO: 1 over the entire length of SEQ ID NO: 1;
- c) a genomic polynucleotide sequence having at least 90% identity to coding regions of SEQ ID NO: 1 over the entire length of SEQ ID NO: 1; and
- d) a genomic polynucleotide sequence having at least 95% identity to coding regions of SEQ ID NO: 1 over the entire length of SEQ ID NO: 1.

11. An intron obtained from a genomic polynucleotide sequence wherein said genomic polynucleotide sequence is selected from the group consisting of:

- a) a genomic polynucleotide sequence having at least 70% identity to coding regions of SEQ ID NO: 10 over the entire length of SEQ ID NO: 10;
- b) a genomic polynucleotide sequence having at least 80% identity to coding regions of SEQ ID NO: 10 over the entire length of SEQ ID NO: 10;
- c) a genomic polynucleotide sequence having at least 90% identity to coding regions of SEQ ID NO: 10 over the entire length of SEQ ID NO: 10; and
- d) a genomic polynucleotide sequence having at least 95% identity to coding regions of SEQ ID NO: 10 over the entire length of SEQ ID NO: 10.

12. A transformed soybean plant comprising a recombinant nucleic acid molecule, the recombinant nucleic acid molecule comprising as operably linked components: (A) a promoter that functions in a plant to cause production of an mRNA molecule; and (B) a nucleic acid sequence that has at least 85% identity to a nucleic acid sequence selected from the group

consisting of SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, complements thereof, and fragments of either.

13. The transformed plant of claim 12, wherein said transformed plant exhibits a reduced palmitic acid level relative to a plant with a similar genetic background but lacking the recombinant nucleic acid molecule.

14. The transformed plant of claim 12, wherein said transformed plant produces a seed with a reduced palmitic acid level relative to a plant with a similar genetic background but lacking the recombinant nucleic acid molecule.

15. The transformed plant of claim 12, wherein said transformed plant exhibits a reduced stearic acid level relative to a plant with a similar genetic background but lacking the recombinant nucleic acid molecule.

16. The transformed plant of claim 12, wherein said transformed plant produces a seed with a reduced stearic acid level relative to a plant with a similar genetic background but lacking the recombinant nucleic acid molecule.

17. The transformed plant of claim 12, wherein said transformed plant produces a seed with a reduced saturated fatty acid content relative to a plant with a similar genetic background but lacking the recombinant nucleic acid molecule.

18. The transformed plant of claim 12, wherein said transformed plant exhibits an increased oleic acid level relative to a plant with a similar genetic background but lacking the recombinant nucleic acid molecule.

19. The transformed plant of claim 12, wherein said transformed plant produces a seed with an increased oleic acid level relative to a plant with a similar genetic background but lacking the recombinant nucleic acid molecule.

20. A transformed soybean plant having a nucleic acid molecule that comprises (a) a first promoter operably linked to a first nucleic acid molecule having a first nucleic acid sequence that has 85% or greater identity to a nucleic acid sequence selected from the group consisting of SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6,


SEQ ID NO: 7, SEQ ID NO: 8, complements thereof, and fragments of either, and (b) a second nucleic acid molecule with a second nucleic acid sequence that encodes an enzyme selected from the group consisting of beta-ketoacyl-ACP synthase I, beta-ketoacyl-ACP synthase IV, and delta-9 desaturase.

21. The transformed soybean plant according to claim 20, wherein the first promoter is a seed specific promoter.


22. The transformed soybean plant according to claim 20, wherein the first promoter is a 7S promoter.

23. The transformed soybean plant according to claim 20, wherein said first nucleic acid molecule is transcribed and is capable of at least partially reducing the level of a transcript encoded by an endogenous *FATB* gene.


24. A method of modifying the lipid composition in a host cell comprising: transforming a host cell with a DNA construct comprising as operably associated components in the 5' to 3' direction of transcription, a transcriptional initiation region functional in said host cell, a DNA sequence selected from the group consisting of SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, complements thereof, and fragments of either, and a transcription termination sequence, and growing said cell under conditions wherein transcription of said DNA sequence is initiated, whereby said lipid composition is modified.




**Figure 1**




**Figure 2**




**Figure 3**




**Figure 4**


**Figure 5**



**Figure 6**



**Figure 7**



**Figure 8**

## SEQUENCE LISTING

<110> Monsanto Technology, LLC

<120> THIOESTERASE-RELATED NUCLEIC ACID SEQUENCES AND METHODS OF USE FOR THE PRODUCTION OF PLANTS WITH MODIFIED FATTY ACID COMPOSITION

<130> 16518.128

<160> 20

<170> PatentIn version 3.1

<210> 1

<211> 4086

<212> DNA

<213> Glycine max

<220>

<223> soybean FATB genomic clone

<400> 1

ttagggaaac aacaaggacg caaatgaca caatagccct ttttcctgt ttccagcttt	60
tctcttctc tctctccatc ttcttcttct tcttcactca gtcaggtagc caaacaatc	120
tgctattcat tcattcattc ctctttctct ctgatcgcaa actgcacctc tacgctccac	180
tcttctcatt ttctcttctt ttctcgcttc tcagatccaa ctcttcagat aacacaagac	240
caaaccgct ttttctgcat ttctagacta gacgttctac cggagaaggt tctcgattct	300
tttctctttt aactttattt ttaaaataat aataatgaga gctggatgag tctgttcgtt	360
gtgaatttgc aggcaatggg gttctcattt tcgttacagt tacagattgc attgtctgct	420
ttctcttctt cccttgcttc ttgctctgt ctgatttttc gtttttattt ctactttta	480
atctttgggg atggatattt tttctgcatt ttttcgggtt gcgatgtttt caggattccg	540
attccgagtc agatctgcgc cggcttatac gacgaatttg ttcttattcg caacttttcg	600
cttgattggc ttgttttacc tctggaatct cacacgtgat caaataagcc tgctatttta	660
gttgaagtag aatttgctt ttatcggaaa gaattctatg gatctgttct gaaattggag	720
ctactgttct gagttgctat tttttttagt agtattaaga acaagtttgc cttttatttt	780
acattttttt cctttgcttt tgccaaaagt ttttatgata actctcttct gtttgtgata	840
taactgatgt gctgtgctgt tattatttgt tatttggggt gaagtataat ttttgggtg	900
aacttgagc atttttagtc cgattgattt ctgatataca tttaaggcta aggttgacct	960
ctaccacgag ttgctgttg atgttttttc catttttttt ttatctcata tcttttacag	1020
tgtttgcta ttgcatctc tcttctttat cccctttctg tggaaagggt ggagggaaaa	1080
tgtatttttt tttctcttc taacttgcgt atattttgca tgcagcgacc ttagaaattc	1140

attatggtgg caacagctgc tacttcatca tttttccctg ttacttcacc ctgcgccggac 1200  
 tctggtggag caggcagcaa acttggtggt gggcctgcaa accttggagg actaaaatcc 1260  
 aaatctgcgt cttctggtgg cttgaaggca aaggcgcaag ccccttcgaa aattaatgga 1320  
 accacagttg ttacatctaa agaaggcttc aagcatgatg atgatctacc ttcgcctccc 1380  
 ccagaaactt ttatcaacca gttgcctgat tggagcatgc ttcttgctgc tatcacaaca 1440  
 attttcttgg ccgctgaaaa gcagtggatg atgcttgatt ggaagccacg gcgacctgac 1500  
 atgcttattg acccctttgg gataggaaaa attgttcagg atggtcttgt gttccgtgaa 1560  
 aacttttcta ttagatcata tgagattggt gctgatcgta ccgcatctat agaaacagta 1620  
 atgaaccatt tgcaagtaag tccgtcctca tacaagtga tctttatgat cttcagagat 1680  
 gagtatgctt tgactaagat agggctgttt atttagacac tgtaattcaa tttcatatat 1740  
 agataatata attctgttgt tacttttcat actatattta tatcaactat ttgcttaaca 1800  
 acaggaaact gcacttaatc atgttaaaag tgctgggctt cttggtgatg gctttggttc 1860  
 cacgccagaa atgtgcaaaa agaacttgat atgggtggtt actcggatgc aggttggtgt 1920  
 ggaacgctat cctacatggt tagtcatcta gattcaacca ttacatgtga tttgcaatgt 1980  
 atccatgtta agctgctatt tctctgtcta ttttagtaat ctttatgagg aatgatcact 2040  
 cctaaatata ttcattgtaa ttattgagac ttaattatga gaaccaaaat gctttggaaa 2100  
 tttgtctggg atgaaaattg attagataca caagctttat acatgatgaa ctatgggaaa 2160  
 ccttggtgcaa cagagctatt gatctgtaca agagatgtag tatagcatta attacatgtt 2220  
 attagataag gtgacttata cttgtttaat tattgtaaaa atagaagctg atactatgta 2280  
 ttctttgcat ttgttttctt accagttata tataccctct gttctgtttg agtactacta 2340  
 gatgtataaa gaatgcaatt attctgactt cttggtgttg ggttgaagtt agataagcta 2400  
 ttagtattat tatggttatt ctaaactcaa ttatctgaaa ttgtgtgtct atatttgctt 2460  
 caggggtgac atagttcaag tggacacttg ggtttctgga tcagggaaga atggtatgcg 2520  
 tcgtgattgg cttttacgtg actgcaaaac tgggtgaaac ttgacaagag cttccaggta 2580  
 gaaatcattc tctgtaattt tccttcccct ttccttctgc ttcaagcaaa ttttaagatg 2640  
 tgtatcttaa tgtgcaagat gctgattgga cacaatttta aatctttcaa acatttaca 2700  
 aagttatgga accctttctt ttctctcttg aagatgcaaa tttgtcacga ctgaagtttg 2760  
 aggaaatcat ttgaattttg caatgttaaa aaagataatg aactacatat tttgcaggca 2820  
 aaaacctcta attgaacaaa ctgaacattg tatcttagtt tatttatcag actttatcat 2880  
 gtgtactgat gcatcacctt ggagcttgta atgaattaca tattagcatt ttctgaactg 2940

tatgttatgg ttttggatg ctacagtgt tgggtcatga tgaataagct gacacggagg 3000  
 ctgtctaaaa ttccagaaga agtcagacag gagataggat cttattttgt ggattctgat 3060  
 ccaattctag aagaggataa cagaaaactg actaaacttg acgacaacac agcggattat 3120  
 attcgtaccg gtttaagtgt atgtcaacta gtttttttgt aattgttgtc attaatttct 3180  
 tttcttaaat tatttcagat gttgctttct aattagttta cattatgtat cttcatttct 3240  
 ccagtcagg tggagtgatc tagatatcaa tcagcatgac aacaatgtga agtacattga 3300  
 ctggattctg gaggtatatt tctgttcttg tattctaata cactgcagtc cttgttttgt 3360  
 tgtaaaccaa aggactgtcc tttgattgtt tgcagagtgc tccacagcca atcttgga 3420  
 gtcagagct ttttccgtg acttttagat ataggagga gtgtgtagg gacagtgtgc 3480  
 tggattccct gactgctgta tctggggccg acatgggcaa tctagctcac agtggacatg 3540  
 ttgagtcaa gcatttgctt cgactcgaaa atgggtgctga gattgtgagg ggcaggactg 3600  
 agtggaggcc caaacctatg aacaacattg gtgttgtaa ccagggtcca gcagaaagca 3660  
 cctaagattt tgaatgggtt aacgggtgga gttgcatcag tctccttgct atgttttagac 3720  
 ttattctggc ctctggggag agttttgctt gtgtctgtcc aatcaatcta catatcttta 3780  
 tatccttcta atttgtgtta ctttgggtgg taagggggaa aagctgcagt aaacctcatt 3840  
 ctctctttct gctgctccat atttcatttc atctctgatt gcgctactgc taggctgtct 3900  
 tcaatattta attgcttgat caaaatagct aggcattgat attattatc ttttctcttg 3960  
 gctcaattaa agatgcaatt ttcatgtga acacagcata actattatc ttattatttt 4020  
 tgtatagcct gtatgcacga atgacttgct catccaatac aaccgtgatt gtatgtccca 4080  
 gctcag 4086

<210> 2  
 <211> 104  
 <212> DNA  
 <213> Glycine max

<220>  
 <223> soybean FATB intron I

<400> 2  
 caaatctgct attcattcat tcattcctct ttctctctga tcgcaaactg cacctctacg 60  
 ctccactctt ctcatcttct cttcctttct cgttctcag atcc 104

<210> 3  
 <211> 839  
 <212> DNA  
 <213> Glycine max

&lt;220&gt;

&lt;223&gt; soybean FATB intron II

&lt;400&gt; 3

ctcgattcctt ttctctttta actttatctt taaaataata ataatgagag ctggatgcgt	60
ctgttcgctt tgaatttcga ggcaatgggg ttctcatttt cgttacagtt acagattgca	120
ttgtctgctt tctctctctc ccttgcttct ttgccttgct tgatttttcg tttttatttc	180
ttacttttaa tttttgggga tggatatttt ttctgcattt tttcgggttg cgatgttttc	240
aggattccga ttccgagtc gatctgcgcc ggcttatacg acgaatttgt tcttattcgc	300
aacttttcgc ttgattggct tgttttacct ctggaatctc acacgtgatc aaataagcct	360
gctatttttag ttgaagtaga atttgttctt tatcggaag aattctatgg atctgttctg	420
aaattggagc tactgtttcg agttgctatt ttttttagta gtattaagaa caagtttgcc	480
ttttatttta catttttttc ctttgctttt gccaaaagt tttatgatca ctctcttctg	540
tttgtgatat aactgatgtg ctgtgctggt attatttggt atttgggtg aagtataatt	600
ttttgggtga acttgagca ttttagtcc gattgatttc tcgatcatc ttaaggctaa	660
ggttgacctc taccacgcgt ttgcgtttga tgtttttcc atttttttt tatctcatat	720
cttttacagt gtttgctat ttgcatttct cttctttatc ccttttctgt ggaaggtggg	780
agggaaaatg tatttttttt ttctcttcta acttgcgtat attttgcag cagcgacct	839

&lt;210&gt; 4

&lt;211&gt; 169

&lt;212&gt; DNA

&lt;213&gt; Glycine max

&lt;220&gt;

&lt;223&gt; soybean FATB intron III

&lt;400&gt; 4

taagtcctgc ctcatacaag tgaatcttta tgatcttcag agatgagat gctttgacta	60
agatagggct gtttatttag acactgtaat tcaatttcac atatagataa tatcattctg	120
ttgttacttt tcatactata tttatatcaa ctatttgctt aacaacagg	169

&lt;210&gt; 5

&lt;211&gt; 525

&lt;212&gt; DNA

&lt;213&gt; Glycine max

&lt;220&gt;

&lt;223&gt; FATB intron IV

&lt;400&gt; 5

gttagtcac tagattcaac cattacatgt gatttgcaat gtatccatgt taagctgcta	60
--	----

```

ttctctgtgc tatttttagta atctttatga ggaatgatca ctctaaata tattcatggt 120
aattattgag acttaattat gagaaccaa atgctttgga aatttgtctg ggatgaaaat 180
tgattagata cacaagcttt atacatgatg aactatggga aaccttgtgc aacagagcta 240
ttgatctgta caagagatgt agtatagcat taattacatg ttattagata aggtgactta 300
tccttgttta attattgtaa aaatagaagc tgatactatg tattctttgc atttgttttc 360
ttaccagtta tatataccct ctgttctgtt tgagtactac tagatgtata aagaatgcaa 420
ttattctgac ttcttggtgt tgggttgaag ttagataagc tattagtatt attatggtta 480
ttctaaatct aattatctga aatttgtgtg ctatatttgc ttcag 525

```

<210> 6  
 <211> 389  
 <212> DNA  
 <213> Glycine max

<220>  
 <223> FATB intron V

```

<400> 6
gtagaaatca ttctctgtaa ttttccttcc cctttccttc tgcttcaagc aaattttaag 60
atgtgtatct taatgtgcac gatgctgatt ggacacaatt ttaaatcttt caaacattta 120
caaaagttat ggaacccttt cttttctctc ttgaagatgc aaatttgtca cgaactgaagt 180
ttgaggaaat catttgaatt ttgcaatgtt aaaaaagata atgaactaca tattttgcag 240
gcaaaaacct ctaattgaac aaactgaaca ttgtatctta gtttatttat cagactttat 300
catgtgtact gatgcatcac cttggagctt gtaatgaatt acatattagc attttctgaa 360
ctgtatgtta tggttttggt gatctacag 389

```

<210> 7  
 <211> 106  
 <212> DNA  
 <213> Glycine max

<220>  
 <223> FATB intron VI

```

<400> 7
tatgtcaact agtttttttg taattgttgt catthaatttc ttttcttaaa ttatttcaga 60
tgttgctttc taattagttt acattatgta tcttcattct tccagt 106

```

<210> 8  
 <211> 82  
 <212> DNA  
 <213> Glycine max

&lt;220&gt;

&lt;223&gt; FATB intron VII

&lt;400&gt; 8

gtatttttct gttcttgtat tctaattccac tgcagtcctt gttttgttgt taaccaaagg 60

actgtccttt gattgtttgc ag 82

&lt;210&gt; 9

&lt;211&gt; 328

&lt;212&gt; PRT

&lt;213&gt; Glycine max

&lt;220&gt;

&lt;223&gt; soybean FATB enzyme

&lt;400&gt; 9

Met Glu Glu Gln Leu Leu Ala Ala Ile Thr Thr Ile Phe Leu Ala Ala  
1 5 10 15Glu Lys Gln Trp Met Met Leu Asp Trp Lys Pro Arg Arg Pro Asp Met  
20 25 30Leu Ile Asp Pro Phe Gly Ile Gly Lys Ile Val Gln Asp Gly Leu Val  
35 40 45Phe Arg Glu Asn Phe Ser Ile Arg Ser Tyr Glu Ile Gly Ala Asp Arg  
50 55 60Thr Ala Ser Ile Glu Thr Val Met Asn His Leu Gln Glu Thr Ala Leu  
65 70 75 80Asn His Val Lys Ser Ala Gly Leu Leu Gly Asp Gly Phe Gly Ser Thr  
85 90 95Pro Glu Met Cys Lys Lys Asn Leu Ile Trp Val Val Thr Arg Met Gln  
100 105 110Val Val Val Glu Arg Tyr Pro Thr Trp Gly Asp Ile Val Gln Val Asp  
115 120 125Thr Trp Val Ser Gly Ser Gly Lys Asn Gly Met Arg Arg Asp Trp Leu  
130 135 140Leu Arg Asp Ser Lys Thr Gly Glu Ile Leu Thr Arg Ala Ser Ser Val  
145 150 155 160Trp Val Met Met Asn Lys Leu Thr Arg Arg Leu Ser Lys Ile Pro Glu  
165 170 175Glu Val Arg Gln Glu Ile Gly Ser Tyr Phe Val Asp Ser Asp Pro Ile  
180 185 190Leu Glu Glu Asp Asn Arg Lys Leu Thr Lys Leu Asp Asp Asn Thr Ala  
195 200 205Asp Tyr Ile Arg Thr Gly Leu Ser Pro Arg Trp Ser Asp Leu Asp Ile  
210 215 220

Asn Gln His Val Asn Asn Val Lys Tyr Ile Gly Trp Ile Leu Glu Ser  
225 230 235 240

Ala Pro Gln Pro Ile Leu Glu Ser His Glu Leu Ser Ser Met Thr Leu  
245 250 255

Glu Tyr Arg Arg Glu Cys Gly Arg Asp Ser Val Leu Asp Ser Leu Thr  
260 265 270

Ala Val Ser Gly Ala Asp Met Gly Asn Leu Ala His Ser Gly His Val  
275 280 285

Glu Cys Lys His Leu Leu Arg Leu Glu Asn Gly Ala Glu Ile Val Arg  
290 295 300

Gly Arg Thr Glu Trp Arg Pro Lys Pro Val Asn Asn Phe Gly Val Val  
305 310 315 320

Asn Gln Val Pro Ala Glu Ser Thr  
325

<210> 10

<211> 1856

<212> DNA

<213> Glycine max

<220>

<223> soybean FATB partial genomic clone

<400> 10

```

ttagggaaac aacaaggacg caaatgaca caatagccct tcttcctgt ttccagcttt      60
tctccttctc tctctccatc ttcttcttct tcttcactca gtcaggtagc caaacaatc     120
tgctattcat tcattcattc ctctttctct ctgatcgcaa actgcacctc tacgctccac     180
tcttctcatt ttctcttctt ttctcgcttc tcagatccaa ctctcagat aacacaagac     240
caaaccgct ttttctgcat ttctagacta gacgttctac cggagaaggt tctcgattct     300
tttctctttt aactttattt ttaaataat aataatgaga gctggatgcg tctgttcgtt     360
gtgaatttcg aggcaatggg gttctcattt tcgttacagt tacagattgc attgtctgct     420
ttcctcttct ccttggttct ttgccttgt ctgatttttc gtttttattt cttactttta     480
atttttgggg atggatattt tttctgcatt ttttcggttt gcgatgtttt caggattccg     540
attccgagtc agatctgcgc cggcttatac gacgaatttg ttctatttcg caacttttcg     600
cttgattggc ttgttttacc tctggaatct cacacgtgat caaataagcc tgctatttta     660
gttgaagtag aatttggtct ttatcggaag gaattctatg gatctgttct gaaattggag     720
ctactgttct gagttgctat tttttttagt agtattaaga acaagtttgc cttttatttt     780
acattttttt cctttgcttt tgccaaaagt ttttatgata actctcttct gtttgtgata     840

```

taactgatgt gctgtgctgt tattatttgt tatttggggt gaagtataat tttttgggtg 900  
 aacttgagc attttagtc cgattgattt ctcgatatca ttttaaggcta aggttgacct 960  
 ctaccacgcg ttgcgtttg atgttttttc cttttttttt ttatctcata ttttttacag 1020  
 tgtttgccta ttgcatctt tcttctttat cccctttctg tggaagggtg gagggaaaat 1080  
 gtattttttt tttctcttct aacttgcgta tattttgcat gcagcgacct tagaaattca 1140  
 ttatggtggc aacagctgct acttcatcat ttttcctgt tacttcaccc tcgccggact 1200  
 ctggtggagc aggcagcaaa cttggtggtg ggctgcaaa ccttgaggga ctaaaatcca 1260  
 aatctgcgtc ttctggtggc ttgaaggcaa aggcgcaagc cccttcgaaa attaattgaa 1320  
 ccacagtgtg tacatctaaa gaaggcttca agcatgatga tgatctacct tcgcctcccc 1380  
 ccagaacttt tatcaaccag ttgcctgatt ggagcatgct tcttgctgct atcacaacaa 1440  
 ttttcttggc cgctgaaaag cagtggatga tgcttgattg gaagccacgg cgacctgaca 1500  
 tgcttattga cccctttggg ataggaaaaa ttgttcagga tggctcttggtg ttccgtgaaa 1560  
 acttttctat tagatcatat gagattggtg ctgatcgtag cgcatctata gaaacagtaa 1620  
 tgaaccattt gcaagtaagt ccgtcctcat acaagtgaat ctttatgatc ttcagagatg 1680  
 agtatgcttt gactaagata gggctgttta tttagacact gtaattcaat ttcatatata 1740  
 gataatatca ttctgttgtt acttttcata ctatatttat atcaactatt tgcttaacaa 1800  
 caggaaactg cacttaatca tgttaaaagt gctgggcttc ttggtgatgg ctggta 1856

<210> 11  
 <211> 34  
 <212> DNA  
 <213> Artificial

<220>  
 <223> Oligonucleotide primer F1

<400> 11 34  
 gcggccgccc cgggttaggg aaacaacaag gacg

<210> 12  
 <211> 34  
 <212> DNA  
 <213> Artificial

<220>  
 <223> Oligonucleotide primer F2

<400> 12 34  
 gcggccgccc cgggcagtca gatccaactc ctca

<210> 13  
<211> 34  
<212> DNA  
<213> Artificial

<220>  
<223> Oligonucleotide primer F3

<400> 13  
gcggccgccc cgggattggt gctgacgta ccgc

34

<210> 14  
<211> 38  
<212> DNA  
<213> Artificial

<220>  
<223> Oligonucleotide primer R1

<400> 14  
gcggccgcgg tccccccct taccaccaa agtatcac

38

<210> 15  
<211> 34  
<212> DNA  
<213> Artificial

<220>  
<223> Oligonucleotide primer R2

<400> 15  
gcggccgcgg taccaaactc tcccaggga acca

34

<210> 16  
<211> 34  
<212> DNA  
<213> Artificial

<220>  
<223> Oligonucleotide primer R3

<400> 16  
gcggccgcgg taccagccat caccaagaag ccca

34

<210> 17  
<211> 37  
<212> DNA  
<213> Artificial

<220>  
<223> Oligonucleotide primer 18133

<400> 17  
gaattcctcg agctcgattc tttctcttt taacttt

37

<210> 18  
 <211> 37  
 <212> DNA  
 <213> Artificial

<220>  
 <223> Oligonucleotide primer 18134

<400> 18  
 gaattcctcg agcatgcaaa atatacgcaa gtagaa

37

<210> 19  
 <211> 854  
 <212> DNA  
 <213> Artificial

<220>  
 <223> PCR product containing soybean FATB intron II

<400> 19  
 gaattcctcg agctcgattc ttttctcttt taactttatt tttaaaataa taataatgag 60  
 agctggatgc gtctgttcgt tgtgaatttc gaggcaatgg gggtctcatt ttcgttacag 120  
 ttacagattg cattgtctgc tttcctcttc tcccttgctt ctttgcttg tctgattttt 180  
 cgtttttatt tcttactttt aatttttggg gatggatatt ttttctgcat tttttcgggt 240  
 tgcgatgttt tcaggattcc gattccgagt cagatctgcg ccggcttata cgacgaattt 300  
 gttcttattc gcaacttttc gcttgattgg cttgttttac ctctggaatc tcacacgtga 360  
 tcaaataagc ctgctatttt agttgaagta gaatttgctt tttatcgga agaattctat 420  
 ggatctgttc tgaaattgga gctactgttt cgagttgcta ttttttttag tagtattaag 480  
 aacaagtttg ccttttattt tacatttttt tcccttgctt ttgccaaaag tttttatgat 540  
 cactctcttc tgtttgtgat ataactgat tgctgtgctg ttattatttg ttatttgggg 600  
 tgaagtataa ttttttgggt gaacttgag catttttagt ccgattgatt tctcgatata 660  
 atttaaggct aaggttgacc tctaccacgc gtttgcgttt gatgtttttt ccattttttt 720  
 tttatctcat atcttttaca gtgtttgctt atttgcatct ctctctctta tcccctttct 780  
 gtggaagggt ggaggggaaaa tgtatttttt ttttctcttc taacttgctt atattttgca 840  
 tgctcgagga attc 854

<210> 20  
 <211> 1688  
 <212> DNA  
 <213> Glycine max

<220>  
 <223> soybean FATB cDNA

<400> 20

acaattacac tgtctctctc ttttccaaaa ttagggaaac aacaaggacg caaatgaca	60
caatagccct tcttccctgt ttccagcttt tctccttctc tctctctcca tcttcttctt	120
cttcttcaact cagtcagatc caactcctca gataacacaa gaccaaaccg gctttttctg	180
catttctaga ctagacgttc taccggagaa gcgaccttag aaattcatta tggtaggcaac	240
agctgctact tcatcatttt tccctgttac ttaccctctg ccggactctg gtggagcagg	300
cagcaaactt ggtggtgggc ctgcaaactt tggaggacta aaatccaaat ctgcgtcttc	360
tggtaggcttg aaggcaaagg cgcaagcccc ttcgaaaatt aatggaacca cagttgttac	420
atctaaagaa agcttcaagc atgatgatga tctaccttcg cctcccccca gaacttttat	480
caaccagttg cctgattgga gcatgcttct tgcgtctatc acaacaattt tcttggccgc	540
tgaaaagcag tggatgatgc ttgattggaa gccacggcga cctgacatgc ttattgaccc	600
ctttgggata ggaaaaattg ttcaggatgg tcttgtgttc cgtgaaaact tttctattag	660
atcatatgag attggtgctg atcgtaccgc atctatagaa acagtaatga accatttgca	720
agaaactgca cttaatcatg ttaaaagtgc tgggcttctt ggtgatggct ttggttccac	780
gccagaaatg tgcaaaaaga acttgatatg ggtggttact cggatgcagg ttgtggtgga	840
acgctatcct acatgggggtg acatagttca agtggacact tgggtttctg gatcagggaa	900
gaatggtatg cgtcgtgatt ggcttttacg tgactccaaa actggtgaaa tcttgacaag	960
agcttccagt gtttgggtca tgatgaataa gctaacacgg aggctgtcta aaattccaga	1020
agaagtcaga caggagatag gatcttattt tgtggattct gatccaattc tggaagagga	1080
taacagaaaa ctgactaaac ttgacgacaa cacagcggat tatattcgta ccggtttaag	1140
tcctagggtg agtgatctag atatcaatca gcatgtcaac aatgtgaagt acattggetg	1200
gattctggag agtgctccac agccaatctt ggagagtcac gagctttctt ccatgacttt	1260
agagtatagg agagagtgtg gtagggacag tgtgctggat tccctgactg ctgtatctgg	1320
ggccgacatg ggcaatctag ctacacagcg gcatgttgag tgcaagcatt tgcttcgact	1380
ggaaaatggt gctgagattg tgaggggcag gactgagtgg aggcccaaac ctgtgaacaa	1440
ctttggtgtt gtgaaccagg ttccagcaga aagcacctaa gatttgaaat ggttaacgat	1500
tggagttgca tcagtctcct tgcctatgtt agacttattc tggttccctg gggagagttt	1560
tgcttggtgc tatccaatca atctacatgt cttaaataat atacaccttc taatttgtga	1620
tactttggtg ggtaaggggg aaaagcagca gtaaatctca ttctcattgt aattaaaaaa	1680
aaaaaaaa	1688